

UNIVERSITY OF SWAZILAND

BACHELOR OF SCIENCE

SUPPLEMENTARY EXAMINATION 2008

TITLE OF PAPER : PHYSICAL CHEMISTRY

COURSE NUMBER : C202

TIME : 3 HOURS

INSTRUCTIONS : THERE ARE SIX QUESTIONS

: ANSWER ANY FOUR QUESTIONS

: BEGIN THE ANSWER TO EACH QUESTION ON
A SEPARATE SHEET OF PAPER

: DATA SHEETS ARE PROVIDED WITH THIS
EXAMINATION PAPER

DO NOT OPEN THIS PAPER UNTIL THE INVIGILATOR INSTRUCTS YOU TO DO
SO.

Question 1(25 marks)

- a) Define the variable, compressibility factor, z . With the aid of Lennard-Jones potential plot, compressibility and isotherm plots, compare and contrast real and ideal gases.

Your account should make mention of interactions, equations and any necessary theories to help clarify your discussion.

[15]

- b) Write short notes **on any One** of the following:

- i) Virial equation [10]
- ii) van der waal's equation [10]

Use diagrams, equations or plots to clarify your notes where necessary.

QUESTION 2 [25 marks]

- a) A real gas equation of state for a gas is given by:

$$P = RT(V_m - \beta)^{-1} - (\alpha/T)V_m^{-2} \quad (1)$$

- (i) Derive an expression for $V_{m,c}$, T_c and P_c . [12]
 - (ii) Find an expression for the Boyle's temperature, T_B . [4]
 - (iii) Estimate the temperature at which oxygen behaves as an ideal gas, T_B given the constants: $\alpha=1.748 \text{ L}^2\text{atm mol}^{-2}\text{K}$ and $\beta=0.0345 \text{ L mol}^{-1}$. [2]
 - (iv) Estimate the radii of real gas molecules using equation (1) for real gases given a critical molar volume of $250 \text{ cm}^3\text{mol}^{-1}$ [4]
- b) Using the critical point expressions for $V_{m,c}$, T_c and P_c find an expression or value for compressibility at the critical point, Z_c [3]

Question 3 [25 Marks]

- a) Derive Kirrchoff's equation: [6]


$$\Delta H_r(T_2) = \Delta H_r(T_1) + \Delta_r C_{p,m} \Delta T$$

where $C_{p,m}$ is temperature independent.

- b) Using the data in the table below calculate

- i) $\Delta_r H^\theta$ at 298 K [4]
- ii) $\Delta_r H$ at 346 K [5]

for the hydrogenation reaction:



	$\text{C}_2\text{H}_4(\text{g})$	$\text{H}_2(\text{g})$	$\text{C}_2\text{H}_2(\text{g})$
$C_{p,m} \text{ J/mol/K}$	43.56	43.93	28.82
$\Delta_f H^\theta \text{ kJ/mol}$	+52.30	0	+226.8

- c) (i) Using an appropriate Master Equation derive the Maxwell's relation

$$(\delta S/\delta V)_T = (\delta P/\delta T)_V \quad [5]$$

(ii) Using the Maxwell's relation in (i) find the expression for internal energy change with volume under isothermal conditions for real gases using Berthelot's relation:


$$(P + an^2/TV^2)(V - nb) = nRT \quad [5]$$

Question 4 [25 Marks]

- a) Using examples and/or diagrams compare and contrast **any one** pair of the following terms
- reversible and irreversible expansion [10]
 - path and state functions [10]
- b) 4 moles of pentane occupies 25 L at 315 K.
- Derive an expression for reversible isothermal expansion. [6]
 - Calculate the work done and heat involved when the gas expands isothermally against a constant external pressure of 115 torr until its volume has doubled. [4]
 - Calculate the efficiency of the system in 1 b (ii) above. [5]

Question 5 [25 Marks]

- a) Write short notes on **any two** of the following
- enthalpy change [5]
 - internal energy change [5]
 - Hess's Law [5]
- b) To Calibrate a calorimeter a 0.120 g naphthalene, C₁₀H₈(s), was burnt at constant volume and it caused the temperature of the calorimeter to rise by 3.05 K. Then 0.10 g of an unknown compound was burned in the same calorimeter, causing a temperature rise of 2.05 K.
- Calculate the heat capacity of the calorimeter [3]
 - Is the unknown compound phenol, C₆H₅OH(s) or ethanol, CH₃CH₂OH(l) whose enthalpies of combustion are Δ_cH⁰ = -3054 kJmol⁻¹ and -1368 kJmol⁻¹ respectively. [4]
- c) Calculate the standard enthalpies of formation of:
- KClO₃(s) from the enthalpy of formation of KCl [4]
 - NOCl(g) from the enthalpy of formation of NO [4]
- Given the attached table and the following information:



Useful information:

	Molecular weights/g mol ⁻¹
Benzoic acid	122.12
D-ribose C ₅ H ₁₀ O ₅ (s)	150.13

QUESTION 6 [25 MARKS]

- a) Write short notes on any Two of the following: [10]
- i) Eutectic temperature and Congruent melting point
 - ii) Zeotrope and Azeotrope
 - iii) Lower consolute and upper consolute temperature
- b) a) Draw a sketch of the phase diagram of carbon dioxide and explain briefly the slopes and curvature of the liquid-solid and the liquid-gas boundaries, respectively. [5]
- c) i) Derive the Clausius-Clapeyron equation for evaporation in the form
- $$\frac{d(\ln p)}{dT} \quad [5]$$
- ii) The triple point of benzene is at 5.5°C and 36 mm Hg. Predict the boiling point of benzene at 0.1 atm pressure. [5]

Useful Relations		General Data						
$(RT)_{298.15K} = 2.4789 \text{ kJ/mol}$		c	$2.997925 \times 10^8 \text{ ms}^{-1}$					
$(RT/F)_{298.15K} = 0.025693 \text{ V}$		e	$1.60219 \times 10^{-19} \text{ C}$					
T/K:	100.15 298.15 500.15 1000.15	F=Le	$9.64846 \times 10^4 \text{ C mol}^{-1}$					
T/Cm⁻¹:	69.61 207.22 347.62 695.13	k	$1.38066 \times 10^{-23} \text{ J K}^{-1}$					
lmmHg=	133.222 N m^{-2}	R=Lk	$8.31441 \text{ J K}^{-1} \text{ mol}^{-1}$					
hc/k=	$1.43878 \times 10^{-2} \text{ m K}$		$8.20575 \times 10^{-2} \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$					
1atm	1 cal	1 eV	1cm⁻¹					
$-1.01325 \times 10^5 \text{ Nm}^{-2}$	-4.184 J	$-1.602189 \times 10^{-19} \text{ J}$	$-0.124 \times 10^{-3} \text{ eV}$					
-760 torr		-96.485 kJ/mol	$-1.9864 \times 10^{-23} \text{ J}$					
-1 bar		$= 8065.5 \text{ cm}^{-1}$						
SI-units:								
$1 \text{ L} = 1000 \text{ ml} = 1000 \text{ cm}^3 = 1 \text{ dm}^3$								
$1 \text{ dm} = 0.1 \text{ m}$								
$1 \text{ cal (thermochemical)} = 4.184 \text{ J}$								
dipole moment: $1 \text{ Debye} = 3.33564 \times 10^{-30} \text{ C m}$								
force: $1 \text{ N} = 1 \text{ J m}^{-1} = 1 \text{ kgms}^{-2} = 10^5 \text{ dyne}$ pressure: $1 \text{ Pa} = 1 \text{ Nm}^{-2} = 1 \text{ Jm}^{-3}$								
$1 \text{ J} = 1 \text{ Nm}$								
power: $1 \text{ W} = 1 \text{ J s}^{-1}$		potential: $1 \text{ V} = 1 \text{ J C}^{-1}$						
magnetic flux: $1 \text{ T} = 1 \text{ Vsm}^{-2} = 1 \text{ JCs}^{-2}$ current: $1 \text{ A} = 1 \text{ Cs}^{-1}$								
Prefixes:								
p	n	m	c	d	k	M	G	
pico	nano	micro	milli	centi	deci	kilo	mega	giga
10^{-12}	10^{-9}	10^{-6}	10^{-3}	10^{-2}	10^{-1}	10^3	10^6	10^9
Bohr radius		a₀						
						$5.29177 \times 10^{-11} \text{ m}$		
Gravitational constant		G						
						$6.67259 \times 10^{-11} \text{ Nm}^2 \text{ kg}^{-2}$		
Gravitational acceleration		g						
						9.80665 ms^{-2}		
Bohr magneton		$\mu_B = \frac{e\hbar}{2m_e}$						
						$9.27402 \times 10^{-24} \text{ JT}^{-1}$		
Nuclear magneton		$\mu_N = \frac{e\hbar}{2m_p}$						
						$5.05079 \times 10^{-27} \text{ JT}^{-1}$		
Planck constant		h						
						$6.62618 \times 10^{-34} \text{ Js}$		
$\hbar = \frac{h}{2\pi}$						$1.05459 \times 10^{-34} \text{ Js}$		
Avogadro constant		L or N_{AV}						
						$6.02214 \times 10^{23} \text{ mol}^{-1}$		
Atomis mass unit		u						
						$1.66054 \times 10^{-27} \text{ kg}$		
Electron mass		m_e						
						$9.10939 \times 10^{-31} \text{ kg}$		
Proton mass		m_p						
						$1.67262 \times 10^{-27} \text{ kg}$		
Neutron mass		m_n						
						$1.67493 \times 10^{-27} \text{ kg}$		
Vacuum permittivity		$\epsilon_0 = \mu_0^{-1} \text{ c}^{-2}$						
						$8.854188 \times 10^{-12} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$		
Vacuum permeability		μ_0						
						$4\pi \times 10^{-7} \text{ Js}^2 \text{ C}^{-2} \text{ m}^{-1}$		
Bohr magneton		$\mu_B = \frac{e\hbar}{2m_e}$						
						$9.27402 \times 10^{-24} \text{ JT}^{-1}$		
Nuclear magneton		$\mu_N = \frac{e\hbar}{2m_p}$						
						$5.05079 \times 10^{-27} \text{ JT}^{-1}$		
Gravitational constant		G						
						$6.67259 \times 10^{-11} \text{ Nm}^2 \text{ kg}^{-2}$		
Gravitational acceleration		g						
						9.80665 ms^{-2}		
Bohr radius		a₀						
						$5.29177 \times 10^{-11} \text{ m}$		

THE PERIODIC TABLE OF ELEMENTS

Group	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18															
	IA	IIA	IIIB	IVB	VB	VIB	VII B	VIII B	VIIIB	IB	IIB	IIIA	IVA	VA	VIA	VIIA	VIIA	VIIIA															
Period 1	1 H 1.008																	2 He 4.003															
2	3 Li 6.94	4 Be 9.01											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18															
3	11 Na 22.99	12 Mg 24.31											13 Al 26.9	14 Si 28.09	15 P 30.97	16 S 32.06	17 Cl 35.45	18 Ar 39.95															
4	19 K 39.10	20 Ca 40.08											21 Sc 44.96	22 Ti 47.90	23 V 50.94	24 Cr 52.01	25 Mn 54.9	26 Fe 55.85	27 Co 58.71	28 Ni 58.71	29 Cu 63.54	30 Zn 65.37	31 Ga 69.7	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.91	36 Kr 83.80					
5	37 Rb 85.47	38 Sr 87.62											39 Y 88.91	40 Zr 91.22	41 Nb 91.22	42 Mo 95.94	43 Tc 98.9	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3					
6	55 Cs 132.9	56 Ba 137.3											71 Lu 174.9	72 Hf 178.5	73 Ta 180.9	74 W 183.8	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 196.9	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 208.9	84 Po 210	85 At 210	86 Rn 222					
7	87 Fr 223	88 Ra 226.0											103 Lr 257	104 Unq	105 Unp	106 Unh	107 Uns	108 Uno	109 Une														

	57	58	59	60	61	62	63	64	65	66	67	68	69	70
Lanthanides	La 138.9	Ce 140.1	Pr 140.9	Nd 144.2	Pm 146.9	Sm 150.9	Eu 151.3	Gd 157.3	Tb 158.9	Dy 162.5	Ho 164.9	Er 167.3	Tm 168.9	Yb 173.0
Actinides	89 Ac 227.0	90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np 237.1	94 Pu 239.1	95 Am 241.1	96 Cm 247.1	97 Bk 249.1	98 Cf 251.1	99 Es 254.1	100 Fm 257.1	101 Md 258.1	102 No 255

Numbers below the symbol indicates the atomic masses; and the numbers above the symbol indicates the atomic numbers.

Standard molar Gibbs free energy and molar entropy of formation at 298.15 K

	M_r	$\Delta G_f^\ominus / \text{kJ/mol}$	$S^\ominus / \text{J K}^{-1} \text{mol}^{-1}$		M_r	$\Delta G_f^\ominus / \text{kJ/mol}$	$S^\ominus / \text{J K}^{-1} \text{mol}^{-1}$		M_r	$\Delta G_f^\ominus / \text{kJ/mol}$	$S^\ominus / \text{J K}^{-1} \text{mol}^{-1}$
$\text{H}_2\text{O}(\text{g})$	18.015	-228.57	188.83	$\text{O}_3(\text{g})$	47.998	163.2	238.93	organic compounds			
$\text{H}_2\text{O}(\text{l})$	18.015	-120.35	109.6	$\text{NO}(\text{g})$	30.006	86.55	210.76	$\text{CH}_4(\text{g})$ methane	16.043	-50.72	186.26
$\text{H}_2\text{O}_2(\text{l})$	34.015	-120.35	109.6	$\text{NO}_2(\text{g})$	46.006	51.31	240.06	$\text{C}_2\text{H}_2(\text{g})$ ethyne	26.038	209.20	200.94
$\text{NH}_3(\text{g})$	17.031	-16.45	192.45	$\text{N}_2\text{O}_4(\text{g})$	92.012	97.89	304.29	$\text{C}_2\text{H}_4(\text{g})$ ethene	28.05	68.15	219.56
$\text{N}_2\text{H}_4(\text{l})$	32.045	149.43	121.21	$\text{SO}_2(\text{g})$	64.063	-300.19	248.22	$\text{C}_2\text{H}_6(\text{g})$ ethane	30.070	-32.82	229.60
$\text{N}_3\text{H}(\text{l})$	43.028	327.3	140.6	$\text{H}_2\text{S}(\text{g})$	34.080	-33.56	205.79	C_3H_6 cyclopropane(g)	42.081	104.45	237.55
$\text{N}_3\text{H}(\text{g})$	43.028	328.1	238.97	$\text{SF}_6(\text{g})$	146.054	-1105.3	291.82	C_3H_6 propene(g)	42.081	62.78	267.05
$\text{HNO}_3(\text{l})$	63.013	-80.71	155.60	$\text{HF}(\text{g})$	20.006	-273.2	173.78	C_4H_{10} n-butane (g)	58.124	-17.03	310.23
$\text{NH}_2\text{OH}(\text{s})$	33.030			$\text{HC}(\text{g})$	36.461	-95.30	186.91	C_5H_{12} n-pentane(g)	72.151	-8.20	348.40
$\text{NH}_4\text{Cl}(\text{s})$	53.492	-202.87	94.6	$\text{HCl}(\text{aq})$	36.461	-131.23	56.5	C_6H_{12} cyclohexane (l)	84.163	26.8	
$\text{HgCl}_2(\text{s})$	271.50	-178.6	146.0	$\text{HBr}(\text{g})$	80.917	-53.45	198.70	C_6H_{14} n-hexane (l)	86.178		204.3
$\text{H}_2\text{SO}_4(\text{l})$	98.078	-690.00	156.90	$\text{HI}(\text{g})$	127.912	1.70	206.59	C_6H_6 benzene (l)	78.115	124.3	173.3
$\text{H}_2\text{SO}_4(\text{aq})$	98.078	-744.53	20.1	$\text{CO}_2(\text{g})$	44.010	-394.36	213.74	C_6H_6 benzene (g)	78.115	129.72	269.31
$\text{NaCl}(\text{s})$	58.443	-384.14	72.13	$\text{CO}(\text{g})$	28.011	-137.17	197.67	C_8H_{18} n-octane (l)	114.233	6.4	361.1
$\text{NaOH}(\text{s})$	39.997	-379.49	64.46	$\text{Al}_2\text{O}_3(\alpha, \text{s})$	101.945	-1582.3	50.92	C_{10}H_8 naphthalene (l)	128.175		
$\text{KCl}(\text{s})$	74.555	-409.14	82.59	SiO_2	60.09	-856.64	41.84	$\text{CH}_3\text{OH}(\text{g})$	32.042	-161.96	239.81
$\text{KBr}(\text{s})$	119.011	-380.66	95.90	$\text{FeS}(\text{s})$	87.91	-100.4	60.29	$\text{CH}_3\text{OH}(\text{l})$	32.042	-166.27	126.8
$\text{KI}(\text{s})$	166.006	-324.89	106.32	$\text{FeS}_2(\text{s})$	119.975	-166.9	52.93	$\text{CH}_3\text{CHO}(\text{g})$	44.054	-128.86	250.3
				$\text{AgCl}(\text{s})$	143.323	-109.79	96.2	$\text{CH}_3\text{CH}_2\text{OH}(\text{l})$	46.07	-174.78	160.7
								$\text{CH}_3\text{COOH}(\text{l})$	60.053	-389.9	159.8
$\text{He}(\text{g})$	4.003	0	126.15	$\text{Hg}(\text{g})$	200.59	31.82	174.96	$\text{CH}_3\text{COOC}_2\text{H}_5(\text{l})$	88.107	-332.7	259.4
$\text{Ar}(\text{g})$	39.95	0	154.84	$\text{Hg}(\text{l})$	200.59	0	76.02	$\text{C}_6\text{H}_5\text{OH}(\text{s})$	94.114	-50.9	146.0
$\text{H}_2(\text{g})$	2.016	0	130.684	$\text{Ag}(\text{g})$	107.87	245.65	173.00	$\text{C}_6\text{H}_5\text{NH}_2(\text{l})$	93.129		
$\text{N}_2(\text{g})$	28.013	0	191.61	$\text{Ag}(\text{s})$	107.87	0	42.55	$\text{CH}_2(\text{NH}_2)\text{CO}_2\text{H}$, glycine (s)	75.068	-373.4	103.5
$\text{O}_2(\text{g})$	31.999	0	205.138	$\text{Na}(\text{g})$	370.95	76.76	153.71	$\text{C}_6\text{H}_{12}\text{O}_6$, α -D-glucose (s)	180.159		
$\text{O}_3(\text{g})$	47.998	163.2	238.93	$\text{Na}(\text{s})$	22.99	0	51.21	$\text{C}_6\text{H}_{22}\text{O}_6$, β -D-glucose (s)	180.159	-910	212
$\text{Cl}_2(\text{g})$	70.91	0	223.07					$\text{C}_{12}\text{H}_{22}\text{O}_{11}$, sucrose (s)	342.303	-1543	360.2
$\text{Br}_2(\text{g})$	159.82	3.110	245.46					$\text{CH}_3\text{CH}(\text{OH})\text{COOH}$	90.079		
$\text{Br}_2(\text{l})$	159.82	0	152.23					lactic acid (s)			
$\text{I}_2(\text{g})$	253.81	19.33	260.69								
$\text{I}_2(\text{s})$	253.81	0	116.135								

Source: American Institute of Physics handbook, McGraw-Hill.

	M_r	$\Delta H_f^\ominus / \text{kJ/mol}$	M_r	$\Delta H_f^\ominus / \text{kJ/mol}$	$a / \text{J K}^{-1} \text{mol}^{-1}$	$b / 10^{-3} \text{J K}^{-2} \text{mol}^{-1}$	$c / 10^5 \text{J K mol}^{-1}$
H ₂ O(g)	18.015	-241.8	O ₃ (g)	47.998	0	0	0
H ₂ O(l)	18.015	-285.8	NO(g)	30.006	+142.7	0	0
H ₂ O ₂ (l)	34.015	-187.8	NO ₂ (g)	46.006	+33.2	3.26	0.50
NH ₃ (g)	17.031	-46.1	N ₂ O(g)	92.012	+9.2	4.18	-1.67
N ₂ H ₄ (l)	32.045	+50.6	SO ₂ (g)	64.063	-296.8	28.58	-0.50
N ₂ H(g)	43.028	+264.1	H ₂ S(g)	34.080	-20.6	3.77	-2.85
N ₃ H(g)	43.028	+294.1	SF ₆ (g)	146.054	-1209	37.03	8.79
HNO ₃ (l)	63.013	-174.1	HF(g)	20.006	-271.1	44.23	-8.62
NH ₂ OH(s)	33.030	-114.2	HCl(g)	36.461	-92.3	30.54	0
NH ₄ Cl(s)	53.492	-314.4	HCl(aq)	36.461	-167.2	25.10	-1.55
HgCl ₂ (s)	271.50	-224.3	HBr(g)	80.917	+36.4	47.86	-1.92
H ₂ SO ₄ (l)	98.078	-814.0	HI(g)	127.912	+26.5		
H ₂ SO ₄ (aq)	98.078	-909.3	CO ₂ (g)	44.010	-393.5		
NaCl(s)	58.443	-411.0	CO(g)	28.011	-110.5		
NaOH(s)	39.997	-426.7	Al ₂ O ₃ (α,s)	101.945	-1675.7		
KCl(s)	74.555	-435.9	SiO ₂ (s)	60.085	-910.9		
KBr(s)	119.011	-392.2	FeS(s)	87.91	-100.0		
KI(s)	166.006	-327.6	FeS ₂ (s)	119.975	-178.2		
Diatomics(g)	—	0	AgCl(s)	143.323	-127.1		
			CH ₄ (g)	16.043	-74.81		
			C ₂ H ₂ (g)	26.038	+226.8		
			C ₂ H ₄ (g)	28.054	+52.30		
			C ₂ H ₆ (g)	30.070	-84.64		
			C ₃ H ₆ cyclopropane(g)	42.081	53.35		
			C ₃ H ₆ propene(g)	42.081	20.5		
			C ₄ H ₁₀ n-butane (g)	58.124	-126.11		
			C ₅ H ₁₂ n-pentane(g)	72.151	-146.4		
			C ₆ H ₁₂ cyclohexane (l)	84.163	-156.2		
			C ₆ H ₁₄ n-hexane (l)	86.178	-198.7		
			C ₆ H ₆ benzene (l)	78.115	+48.99		
			C ₈ H ₁₈ n-octane (l)	114.233	-249.8		
			C ₁₀ H ₈ naphthalene (l)	128.175	+78.53		
			CH ₃ OH (l)	32.042	-239.0		
			CH ₃ CHO (g)	44.054	-166.0		
			CH ₃ CH ₂ OH (l)	46.070	-277.0		
			CH ₃ COOH (l)	60.053	-484.2		
			CH ₃ COOC ₂ H ₅ (l)	88.107	-486.6		
			C ₆ H ₅ OH (s)	94.114	-165.0		
			C ₆ H ₅ NH ₂ (l)	93.129	-31.1		
			NH ₂ CO.NH ₂ urea(s)	60.056	-333.0		
			CH ₂ (NH ₂)CO ₂ H, glycine (s)	75.068	-537.2		
			C ₆ H ₁₂ O ₆ , α-D-glucose (s)	180.159	-1274		
			C ₆ H ₁₂ O ₆ , β-D-glucose (s)	180.159	-1268		
			C ₁₂ H ₂₂ O ₁₁ , sucrose (s)	342.303	-2222		
			CH ₃ CH(OH)COOH lactic acid (s)	90.079	-694.0		

l. Sublimation; ^a various pressures; ^b at 1 atm

Standard molar Gibbs free energy and molar entropy of formation at 298.15 K

M _r	ΔG _f ⁰ /kJ/mol	S ⁰ /J K ⁻¹ mol ⁻¹	M _r	ΔG _f ⁰ /kJ/mol	S ⁰ /J K ⁻¹ mol ⁻¹	M _r	ΔG _f ⁰ /kJ/mol	S ⁰ /J K ⁻¹ mol ⁻¹
H ₂ O(g)	18.015	-228.57	188.83	O ₃ (g)	163.2	47.998	163.2	238.93
H ₂ O(l)	18.015	-120.35	109.6	NO(g)	86.55	30.006	86.55	210.76
H ₂ O ₂ (l)	34.015	-120.35	109.6	NO ₂ (g)	51.31	46.006	51.31	240.06
NH ₃ (g)	17.031	-16.45	192.45	N ₂ O ₄ (g)	97.89	92.012	97.89	304.29
N ₂ H ₄ (l)	32.045	149.43	121.21	SO ₂ (g)	-300.19	64.063	-300.19	248.22
N ₃ H(l)	43.028	327.3	140.6	H ₂ S(g)	-33.56	34.080	-33.56	205.79
N ₃ H(g)	43.028	328.1	238.97	SF ₆ (g)	-1105.3	146.054	-1105.3	291.82
HNHO ₃ (l)	63.013	-80.71	155.60	HF(g)	-273.2	20.006	-273.2	173.78
NH ₂ OH(s)	33.030			HCl(g)	-95.30	36.461	-95.30	186.91
NH ₄ Cl(s)	53.492	-202.87	94.6	HCl(aq)	-131.23	36.461	-131.23	56.5
HgCl ₂ (s)	271.50	-178.6	146.0	HBr(g)	-53.45	80.917	-53.45	198.70
H ₂ SO ₄ (l)	98.078	-690.00	156.90	HI(g)	1.70	127.912	1.70	206.59
H ₂ SO ₄ (aq)	98.078	-744.53	20.1	CO ₂ (g)	-394.36	44.010	-394.36	213.74
NaCl(s)	58.443	-384.14	72.13	CO(g)	-137.17	28.011	-137.17	197.67
NaOH(s)	39.997	-379.49	64.46	AL ₂ O ₃ (α,s)	-1582.3	101.945	-1582.3	50.92
KCl(s)	74.555	-409.14	82.59	SiO ₂	-856.64	60.09	-856.64	41.84
KBr(s)	119.011	-380.66	95.90	FeS(s)	-100.4	87.91	-100.4	60.29
KI(s)	166.006	-324.89	106.32	FeS ₂ (s)	-166.9	119.975	-166.9	52.93
				AgCl(s)	-109.79	143.323	-109.79	96.2
He(g)	4.003	0	126.15	Hg(g)	31.82	200.59	31.82	174.96
Ar(g)	39.95	0	154.84	Hg(l)	0	200.59	0	76.02
H ₂ (g)	2.016	0	130.684	Ag(g)	245.65	107.87	245.65	173.00
N ₂ (g)	28.013	0	191.61	Ag(s)	0	107.87	0	42.55
O ₂ (g)	31.999	0	205.138	Na(g)	76.76	370.95	76.76	153.71
O ₃ (g)	47.998	163.2	238.93	Na(s)	0	22.99	0	51.21
Cl ₂ (g)	70.91	0	223.07					
Br ₂ (g)	159.82	3.110	245.46					
Br ₂ (l)	159.82	0	152.23					
I ₂ (g)	253.81	19.33	260.69					
I ₂ (s)	253.81	0	116.135					

organic compounds	M _r	ΔG _f ⁰ /kJ/mol	S ⁰ /J K ⁻¹ mol ⁻¹
CH ₄ (g) methane	16.043	-50.72	186.26
C ₂ H ₂ (g) ethyne	26.038	209.20	200.94
C ₂ H ₄ (g) ethene	28.05	68.15	219.56
C ₂ H ₆ (g) ethane	30.070	-32.82	229.60
C ₃ H ₆ cyclopropane(g)	42.081	104.45	237.55
C ₃ H ₆ propene(g)	42.081	62.78	267.05
C ₄ H ₁₀ n-butane (g)	58.124	-17.03	310.23
C ₅ H ₁₂ n-pentane(g)	72.151	-6.20	348.40
C ₆ H ₁₂ cyclohexane (l)	84.163	26.8	
C ₆ H ₁₄ n-hexane (l)	86.178		204.3
C ₆ H ₆ benzene (l)	78.115	124.3	173.3
C ₆ H ₆ benzene (g)	78.115	129.72	269.31
C ₈ H ₁₈ n-octane (l)	114.233	6.4	361.1
C ₁₀ H ₈ naphthalene (l)	128.175		
CH ₃ OH (g)	32.042	-161.96	239.81
CH ₃ OH (l)	32.042	-166.27	126.8
CH ₃ CHO (g)	44.054	-128.86	250.3
CH ₃ CH ₂ OH (l)	46.07	-174.78	160.7
CH ₃ COOH (l)	60.053	-389.9	159.8
CH ₃ COOC ₂ H ₅ (l)	88.107	-332.7	259.4
C ₆ H ₅ OH (s)	94.114	-50.9	146.0
C ₆ H ₅ Ni ₂ (l)	93.129		
CH ₂ (NH ₂)CO ₂ H, glycine (s)	75.068	-373.4	103.5
C ₆ H ₁₂ O ₆ , α-D-glucose (s)	180.159		
C ₆ H ₁₂ O ₆ , β-D-glucose (s)	180.159	-910	212
C ₁₂ H ₂₂ O ₁₁ , sucrose (s)	342.303	-1543	360.2
CH ₃ CH(OH)COOH	90.079		
lactic acid (s)			

Source: American Institute of Physics handbook, McGraw-Hill.

Heat capacities at 25°C

	$C_{v,m}$	$C_{p,m}$
	$\text{JK}^{-1} \text{mol}^{-1}$	$\text{JK}^{-1} \text{mol}^{-1}$
He, Ne, Ar, Kr, Xe	12.47	20.78
H ₂	20.50	28.81
O ₂	21.01	29.33
N ₂	20.83	29.14
CO ₂	28.83	37.14
NH ₃	27.17	35.48
CH ₄	27.43	35.74

F.P. Depression, B.P. Elevation

Solvent	F.P. °C	K_f °C kg mol ⁻¹	B.P. (°C, 101kNm ⁻²)	K_b °C kg mol ⁻¹
Water	0	1.86	100.0	0.52
Benzene	5.51	5.10	80.1	2.60
Acetic Acid	16.6	3.90	118.1	3.10
Cyclohexane	6.5	20.2	81.4	2.79
Camphor	177.7	40.0	205	-
Nitrobenzene	5.7	6.9	210.9	5.24
Ethanol	-177	-	78.5	1.22
Chloroform	-64	-	61.3	3.63

Third Law entropies at 25°C, $\text{Sm}^{\circ}/\text{J K}^{-1} \text{mol}^{-1}$

	Solids	Liquids	Gases
Ag	42.68	Hg	H ₂
C(gt)	5.77	Br ₂	N ₂
C(d)	2.44		O ₂
Cu	33.4		Cl ₂
Zn	41.6	H ₂ O	
I ₂	116.7		CO ₂
S(Rh)	31.9	HNO ₃	HCl
			H ₂ S
AgCl	96.2	C ₂ H ₅ OH	NH ₃
AgBr	104.6	CH ₃ OH	CH ₄
CuSO ₄ ·5H ₂ O	305.4	C ₆ H ₆	C ₂ H ₆
HgCl ₂	144	CH ₃ COOH	CH ₃ CHO
Sucrose	360.2	C ₆ H ₁₂	