

UNIVERSITY OF SWAZILAND
SUPPLEMENTARY EXAMINATION 2008

TITLE OF PAPER: PHYSICAL CHEMISTRY

COURSE NUMBER: C302

TIME: THREE (3) HOURS

INSTRUCTIONS:

There are **six** questions. Each question is worth 25 marks. Answer **any four** questions.

A list of integrals, a data sheet and a periodic table are attached

Non-programmable electronic calculators may be used.

DO NOT OPEN THIS PAPER UNTIL PERMISSION TO DO SO HAS BEEN GRANTED BY THE CHIEF INVIGILATOR.

Question 1(25marks)

- a. Explain how Einstein's introduction of quantization accounted for the heat capacities of metals at low temperatures. [5]
- b. The work function of platinum is 5.65 eV.
- What is the minimum frequency of light required to observe the photoelectric effect on Pt? [3]
 - If light with 150 nm wavelength is absorbed by the surface, what is the velocity of the ejected electrons? [3]
 - What is the wavelength associated with the electron traveling at this speed? [3]
- c. Show that $f(x) = e^{2x}$ is an eigenfunction of the operator $\hat{\Omega} = \frac{d^2}{dx^2} + 2\frac{d}{dx} + 3$ and determine the eigen value. [5]
- d. Determine the commutator of the operators $\hat{A} = x^2 \frac{d}{dx}$ and $\hat{B} = \frac{d^2}{dx^2}$ [6]

Question 2 (25 marks)

- a. Consider the function $\psi(x) = c\sin\pi x/a + d\sin 2\pi x/a$.
- Is $\psi(x)$ an acceptable function for the particle in a one dimensional box of length a ? [4]
 - Is $\psi(x)$ an eigenfunction of the total energy operator $\hat{H} = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2}$? [4]
- b. Consider a particle confined to a one-dimensional box of length L , and whose wavefunction is $\psi(x) = \sqrt{\frac{2}{L}} \sin \frac{n\pi x}{L}$, $n = 1, 2, 3, \dots$
- What are the most likely locations of the particle when $n = 3$? [5]
 - Calculate the probability that the particle will be found between $0.49L$ and $0.51L$ when $n=1$ and when $n=2$. [4]
- c. A two dimensional oscillator has the potential energy
- $$V = \frac{1}{2}k(x^2 + y^2)$$
- Write down the expression for the Schrödinger equation for this system. [3]
 - The energy levels of a one dimensional oscillator are $E = (v + \frac{1}{2})h\nu$, $v=0, 1, 2, \dots$. Use this to write an expression for the energy levels of a two dimensional oscillator. [2]
 - What is the degeneracy of the first four energy levels. [3]

Question 3 (25 marks)

- a. One of the excited states of the hydrogen atom is described by the wavefunction

$$\psi = \left(2 - \frac{r}{a_0}\right) e^{-r/2a_0}$$

- (i) Normalize ψ to 1. [6]
- (ii) Evaluate the expectation value of r for the hydrogen atom with the above wavefunction. [7]
- b. Specify and account for the selection rules for transitions in hydrogenic atoms. [4]
- c. What atomic terms are possible for the electron configuration ns^1np^1 ? Which term is likely to lie lowest in energy? [5]
- d. What values of J may occur in the term 3D . How many states (distinguished by the quantum number M_J) belong to each level? [3]

Question 4 (25 marks)

- a. Distinguish between a bonding and an anti-bonding orbital. [5]
- b. Use molecular orbital theory to explain why the binding energy of N_2^+ is less than that of N_2 whilst that of O_2^+ is greater than that of O_2 . [6]
- c. The ground state of O_2^+ is $^2\Pi_g$ and the next few excited states are $^4\Pi_u$, $^2\Pi_u$, $^4\Sigma_g^-$, $^2\Delta_g$, $^2\Sigma_g^-$, and $^4\Sigma_u^-$. On the basis of the selection rules, which of the excited states can be accessed from the ground state by absorption of UV light? [6]
- d. Predict the number of unpaired electrons and the ground state term of (i) NO and (ii) CO [8]

Question 5(25 marks)

- a. The following lines were observed in the microwave spectrum of H^{127}I and D^{127}I between 60 cm^{-1} and 90 cm^{-1} :

Molecule	Frequency, cm^{-1}			
H^{127}I	64.275	77.130	89.985	
D^{127}I	65.070	71.577	78.084	84.591

- (i) Calculate the values of B , I , and r_e for each molecule. [12]
(ii) In each case what transition gives rise to the first line above?
[4]
(atomic masses are H 1.0078 u, D 2.0140 u and ^{127}I 126.9045 u)
- b. Write down the expression for vibrational-rotational energy levels of a diatomic molecule assumed to be rigid. [3]
- c. Obtain a general expression for the change in energy of the R-branch in HCl in the lowest vibrational state. [4]

Question 6 (25 marks)

- a. The fundamental and first overtone transitions of $^{12}\text{C}^{16}\text{O}$ occur at 2143.0 cm^{-1} and 4260.0 cm^{-1} , respectively. Given that the isotopic masses of ^{12}C and ^{16}O are 12 u (exactly) and 15.9949 u, respectively, calculate
- (i) The equilibrium vibrational frequency [5]
(ii) The anharmonicity constant [3]
(iii) The exact zero point energy [3]
(iv) The force constant of the molecule [4]
- b. The N_2O molecule has three strong bands in its infrared spectrum at 588.8 cm^{-1} , 1285.0 cm^{-1} , and 2223.5 cm^{-1} . All have been shown to be fundamentals and the molecule has been shown to be linear.
- (i) Explain why CO_2 , which is also linear, has only two fundamental IR bands while N_2O has three. [5]
(ii) Where would you look for the overtone and combination bands in the IR spectrum of N_2O ? [5]

Useful Integrals and relations

$$d\tau = r^2 dr \sin\theta d\theta d\phi$$

$$\int x^n \exp(-ax) dx = \frac{n!}{a^{n+1}} \quad (a > 0, n \text{ positive integer})$$

$$\int \sin x dx = -\cos x$$

$$\int \sin^2 ax dx = \frac{x}{2} - \frac{1}{4a} \sin 2ax$$

$$2 \sin\theta \cos\theta = \sin 2\theta$$

General data and fundamental constants

Quantity	Symbol	Value
Speed of light	c	$2.997\,924\,58 \times 10^8 \text{ m s}^{-1}$
Elementary charge	e	$1.602\,177 \times 10^{-19} \text{ C}$
Faraday constant	$F = N_A e$	$9.6485 \times 10^4 \text{ C mol}^{-1}$
Boltzmann constant	k	$1.380\,66 \times 10^{-23} \text{ J K}^{-1}$
Gas constant	$R = N_A k$	$8.314\,51 \text{ J K}^{-1} \text{ mol}^{-1}$ $8.205\,78 \times 10^{-2} \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$ $6.2364 \times 10 \text{ L Torr K}^{-1} \text{ mol}^{-1}$
Planck constant	h	$6.626\,08 \times 10^{-34} \text{ J s}$
	$\hbar = h/2\pi$	$1.054\,57 \times 10^{-34} \text{ J s}$
Avogadro constant	N_A	$6.022\,14 \times 10^{23} \text{ mol}^{-1}$
Atomic mass unit	u	$1.660\,54 \times 10^{-27} \text{ Kg}$
Mass		
electron	m_e	$9.109\,39 \times 10^{-31} \text{ Kg}$
proton	m_p	$1.672\,62 \times 10^{-27} \text{ Kg}$
neutron	m_n	$1.674\,93 \times 10^{-27} \text{ Kg}$
Vacuum permittivity	$\epsilon_0 = 1/c^2 \mu_0$	$8.854\,19 \times 10^{-12} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$
	$4\pi\epsilon_0$	$1.112\,65 \times 10^{-10} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$
Vacuum permeability	μ_0	$4\pi \times 10^{-7} \text{ J s}^2 \text{ C}^{-2} \text{ m}^{-1}$ $4\pi \times 10^{-7} \text{ T}^2 \text{ J}^{-1} \text{ m}^3$
Magneton		
Bohr	$\mu_B = e\hbar/2m_e$	$9.274\,02 \times 10^{-24} \text{ J T}^{-1}$
nuclear	$\mu_N = e\hbar/2m_p$	$5.050\,79 \times 10^{-27} \text{ J T}^{-1}$
g value	g_e	2.002 32
Bohr radius	$a_0 = 4\pi\epsilon_0 \hbar^2 / m_e e^2$	$5.291\,77 \times 10^{-11} \text{ m}$
Fine-structure constant	$\alpha = \mu_0 e^2 c / 2h$	$7.297\,35 \times 10^{-3}$
Rydberg constant	$R_\infty = m_e e^4 / 8h^3 c \epsilon_0^2$	$1.097\,37 \times 10^7 \text{ m}^{-1}$
Standard acceleration of free fall	g	$9.806\,65 \text{ m s}^{-2}$
Gravitational constant	G	$6.672\,59 \times 10^{-11} \text{ N m}^2 \text{ Kg}^{-2}$

Conversion factors

1 cal =	4.184 joules (J)	1 erg =	$1 \times 10^{-7} \text{ J}$
1 eV =	$1.602\,2 \times 10^{-19} \text{ J}$	1 eV/molecule =	96 485 kJ mol ⁻¹

Prefixes	f	p	n	μ	m	c	d	k	M	G
	femto	pico	nano	micro	milli	centi	deci	kilo	mega	giga
	10^{-15}	10^{-12}	10^{-9}	10^{-6}	10^{-3}	10^{-2}	10^{-1}	10^3	10^6	10^9

PERIODIC TABLE OF ELEMENTS

GROUPS

PERIODS	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
	IA	IIA	IIIB	IVB	VB	VIB	VIIIB	VIIIB	VIIIB	IB	IIB	IIIA	IIIA	IVA	VA	VIA	VIIA	VIIIA	
1	1.008 H 1																		4.003 He 2
2	6.941 Li 3	9.012 Be 4											10.811 B 5	12.011 C 6	14.007 N 7	15.999 O 8	18.998 F 9	20.180 Ne 10	
3	22.990 Na 11	24.305 Mg 12											26.982 Al 13	28.086 Si 14	30.974 P 15	32.06 S 16	35.453 Cl 17	39.948 Ar 18	
4	39.098 K 19	40.078 Ca 20	44.956 Sc 21	47.88 Ti 22	50.942 V 23	51.996 Cr 24	54.938 Mn 25	55.847 Fe 26	58.933 Co 27	58.69 Ni 28	63.546 Cu 29	65.39 Zn 30	69.723 Ga 31	72.61 Ge 32	74.922 As 33	78.96 Se 34	79.904 Br 35	83.80 Kr 36	
5	85.468 Rb 37	87.62 Sr 38	88.906 Y 39	91.224 Zr 40	92.906 Nb 41	95.94 Mo 42	98.907 Tc 43	101.07 Ru 44	102.91 Rh 45	106.42 Pd 46	107.87 Ag 47	112.41 Cd 48	114.82 In 49	118.71 Sn 50	121.75 Sb 51	127.60 Te 52	126.90 I 53	131.29 Xe 54	
6	132.91 Cs 55	137.33 Ba 56	138.91 *La 57	178.49 Hf 72	180.95 Ta 73	183.85 W 74	186.21 Re 75	190.2 Os 76	192.22 Ir 77	195.08 Pt 78	196.97 Au 79	200.59 Hg 80	204.38 Tl 81	207.2 Pb 82	208.98 Bi 83	(209) Po 84	(210) At 85	(222) Rn 86	
7	223 Fr 87	226.03 Ra 88	(227) **Ac 89	(261) Rf 104	(262) Ha 105	(263) Unh 106	(262) Uns 107	(265) Uno 108	(266) Une 109	(267) Uun 110									

TRANSITION ELEMENTS

Atomic mass →
Symbol →
Atomic No. →

* Lanthanide Series

** Actinide Series

140.12 Ce 58	140.91 Pr 59	144.24 Nd 60	(145) Pm 61	150.36 Sm 62	151.96 Eu 63	157.25 Gd 64	158.93 Tb 65	162.50 Dy 66	164.93 Ho 67	167.26 Er 68	168.93 Tm 69	173.04 Yb 70	174.97 Lu 71
232.04 Th 90	231.04 Pa 91	238.03 U 92	237.05 Np 93	(244) Pu 94	(243) Am 95	(247) Cm 96	(247) Bk 97	(251) Cf 98	(252) Es 99	(257) Fm 100	(258) Md 101	(259) No 102	(260) Lr 103

() indicates the mass number of the isotope with the longest half-life.