

UNIVERSITY OF SWAZILAND

BACHELOR OF SCIENCE

SUPPLEMENTARY EXAMINATION 2012

TITLE OF PAPER : INTRODUCTORY PHYSICAL CHEMISTRY

COURSE NUMBER : C202

TIME : 3 HOURS

INSTRUCTIONS : THERE ARE SIX QUESTIONS

: ANSWER ANY FOUR QUESTIONS

: BEGIN THE ANSWER TO EACH QUESTION ON
A SEPARATE SHEET OF PAPER

: DATA SHEETS ARE PROVIDED WITH THIS
EXAMINATION PAPER

DO NOT OPEN THIS PAPER UNTIL THE INVIGILATOR INSTRUCTS YOU TO DO
SO.

Question 1 [25 marks]

The compressibility factor, Z , for a real gas is given by

$$Z = PV / nRT$$

and is a measure of the ideal behaviour of the gas.

(a) Use the following data to plot Z versus P for O_2 at 273 K.

$P/(\text{atm})$	1	100	200	300	500	700	900
$V_m/(\text{L}\cdot\text{mol}^{-1})$	22.4138	0.2077	0.1024	0.0719	0.0518	0.0444	0.0403

Where V_m is the molar volume.

[10]

(b) Using the data in "a" compare and contrast real gases and ideal gases. [15]

Your account should make mention of interactions, equations and any necessary theories to help clarify your discussion. You may use additional diagrams and examples to clarify your points.

QUESTION 2 [25 marks]

a) Give a comparison between adiabatic expansion and isothermal expansion. [10]

You may use a graph or equations to illustrate your point.

b) Derive the expression for the change in temperature of an adiabatic expansion of an ideal gas against constant external pressure from V_i to V_f . [5]

c) State and explain the equipartition principle. Using CO as an example show how the principle could be used to evaluate the heat capacity ratio: $\gamma = C_{p,m}/C_{v,m}$ [5]



d) A sample of Krypton, Kr, at 1.5 atm pressure and 298 K expands reversibly and adiabatically from 0.50 L to 1.5 L. Given $C_{v,m} = 12.48 \text{ J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$: Calculate:

- i) the final pressure [3]
- ii) the final temperature [2]

Question 3 [25 Marks]

a) Compare and contrast between enthalpy and internal energy change [5]


b) Find $\Delta_r H^\theta$ for the following reactions from standard enthalpies of formation:



iii)

c) Calculate the standard enthalpies of formation of:

$\text{KClO}_3(\text{s})$ from the enthalpy of formation of KCl [6]



d) Derive Kirrchoff's equation [2]

$$\Delta_r H(T_2) = \Delta_r H(T_1) + \Delta_r C_{p,m} \Delta T$$

Using the data in the table below calculate $\Delta_r H$ and $\Delta_r U$ for the reaction:



At 348 K [5]

	C(graphite)	H ₂ O(g)	CO(g)	H ₂ (g)
$C_{p,m} \text{ J mol}^{-1} \text{ K}^{-1}$	8.53	33.58	29.14	28.82
$\Delta_f H/\text{kJ/mol}$	0	-241.8	-110.5	0

e) To Calibrate a calorimeter a sample of D-ribose ($\text{C}_5\text{H}_{10}\text{O}_5$) of mass 0.727 was burned at constant volume and it caused the temperature of the calorimeter to rise by 0.910 K. Then 0.825 g of an unknown compound was burned in the same calorimeter, causing a temperature rise of 1.94 K.

i) Calculate the heat capacity of the calorimeter [2]

ii) Is the unknown compound phenol, naphthalene C_{10}H_8 (l) or sucrose, $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ (s) whose enthalpies of combustion are $\Delta_c H^\theta = -5157 \text{ kJ mol}^{-1}$ and $-5645 \text{ kJ mol}^{-1}$, respectively. [3]

Question 4 [25 Marks]

a) Briefly discuss the statistical view of entropy [10]

b) 1.00 mol of perfect gas at 27°C is expanded isothermally from an initial pressure of 3.00 atm to a final pressure of 1.00 atm. Calculate q , w , ΔS_{sys} , ΔS_{surr} and ΔS_{tot} if the expansion is done:
(1) reversibly, and [5]

- (2) against a constant external pressure of 1.00 atm. [5]
 (3) adiabatically against a constant pressure of 1.00 atm. [5]


Question 5 [25 Marks]

- a) Derive the integrated Gibbs-Helmholtz equation [5]

$$\frac{\Delta G_2}{T_2} - \frac{\Delta G_1}{T_1} = \Delta H \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

from the fundamental thermodynamic equation of dG

- b) Given the reaction:



Calculate the change in Gibbs free energy ΔG^θ

- i) at 298 K [5]
 ii) at 375 K [5]
 iii) Comment on the significance of the values obtained in (i) and (ii). [2]
 iv) Using the appropriate data and thermodynamic expression, calculate the maximum expansion work for the reaction at 298 K. [3]

- c) The Master Equations states that $dU=TdS-PdV$ and $dA=-PdV-SdT$.

- (i) Using the appropriate Master Equation above derive the Maxwell's relation

$$(\delta S/\delta V)_T = (\delta P/\delta T)_V \quad [3]$$

- (ii) Using the Maxwell's relation in (i) find the expression for internal energy change with volume under isothermal conditions for real gases using Berthelot's relation:

$$(P+an^2/TV^2)(V-nb)=nRT \quad [2]$$

QUESTION 6 [25 MARKS]

- a) Draw a sketch of the phase diagram of carbon dioxide and explain briefly the slopes and curvature of the liquid-solid and the liquid-gas boundaries, respectively. [10]
- b) i) Derive the Clausius-Clapeyron equation for evaporation. [5]
 ii) The triple point of benzene is at 5.5°C and 36 mm Hg. Predict the boiling point of benzene at 0.2 atm pressure. [5]

- c) Given initial concentration of acetic acid, CH_3COOH of 0.25 M calculate the equilibrium concentrations of all ionic species given that the dissociation constant is 1.75×10^{-5} . [5]

END OF EXAM

THE PERIODIC TABLE OF ELEMENTS

Group	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
	IA	IIA	IIIB	IVB	VB	VIB	VII B	VIII B			IB	II B	IIIA	IVA	VA	VIA	VIIA	VIIIA
Period 1	1 H 1.008	NON-METALS ←																2 He 4.003
2	3 Li 6.94	4 Be 9.01	METALLOIDS ←										5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18
3	11 Na 22.99	12 Mg 24.31	METALS →										13 Al 26.9	14 Si 28.09	15 P 30.97	16 S 32.06	17 Cl 35.45	18 Ar 39.95
4	19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.90	23 V 50.94	24 Cr 52.01	25 Mn 54.9	26 Fe 55.85	27 Co 58.71	28 Ni 58.71	29 Cu 63.54	30 Zn 65.37	31 Ga 69.7	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.91	36 Kr 83.80
5	37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 91.22	42 Mo 95.94	43 Tc 98.9	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3
6	55 Cs 132.9	56 Ba 137.3	71 Lu 174.9	72 Hf 178.5	73 Ta 180.9	74 W 183.8	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 196.9	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 208.9	84 Po 210	85 At 210	86 Rn 222
7	87 Fr 223	88 Ra 226.0	103 Lr 257	104 Unq	105 Unp	106 Unh	107 Uns	108 Uno	109 Une									

Lanthanides	57 La 138.9	58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm 146.9	62 Sm 150.9	63 Eu 151.3	64 Gd 157.3	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0
Actinides	89 Ac 227.0	90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np 237.1	94 Pu 239.1	95 Am 241.1	96 Cm 247.1	97 Bk 249.1	98 Cf 251.1	99 Es 254.1	100 Fm 257.1	101 Md 258.1	102 No 255

Numbers below the symbol indicates the atomic masses; and the numbers above the symbol indicates the atomic numbers.

SOURCE: International Union of Pure and Applied Chemistry, 1 mills, ed., Quantities, Units, and symbols in Physical Chemistry, Blackwell Scientific publications, Boston.

Useful Relations				General Data							
$(RT)_{298-15K}=2.4789 \text{ kJ/mol}$				speed of light	c	$2.997\ 925 \times 10^8 \text{ ms}^{-1}$					
$(RT/F)_{298-15K}=0.025\ 693 \text{ V}$				charge of proton	e	$1.602\ 19 \times 10^{-19} \text{ C}$					
T/K: 100.15 298.15 500.15 1000.15				Faraday constant	$F=Le$	$9.648\ 46 \times 10^4 \text{ C mol}^{-1}$					
T/Cm ⁻¹ : 69.61 207.22 347.62 695.13				Boltzmann constant	k	$1.380\ 66 \times 10^{-23} \text{ J K}^{-1}$					
1mmHg=133.222 N m ⁻²				Gas constant	$R=Lk$	$8.314\ 41 \text{ J K}^{-1} \text{ mol}^{-1}$					
hc/k=1.438 78x10 ⁻² m K						$8.205\ 75 \times 10^{-2} \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$					
1atm	1 cal	1 eV	1cm ⁻¹								
$-1.01325 \times 10^5 \text{ Nm}^{-2}$	=4.184 J	=1.602 189x10 ⁻¹⁹ J	=0.124x10 ⁻³ eV	Planck constant	h	$6.626\ 18 \times 10^{-34} \text{ Js}$					
-760torr		=96.485 kJ/mol	=1.9864x10 ⁻²³ J		$\hbar = \frac{h}{2\pi}$	$1.054\ 59 \times 10^{-34} \text{ Js}$					
-1 bar		= 8065.5 cm ⁻¹		Avogadro constant	$L \text{ or } N_{AV}$	$6.022\ 14 \times 10^{23} \text{ mol}^{-1}$					
SI-units:				Atomis mass unit	u	$1.660\ 54 \times 10^{-27} \text{ kg}$					
$1 \text{ L} = 1000 \text{ ml} = 1000 \text{ cm}^3 = 1 \text{ dm}^3$				Electron mass	m_e	$9.109\ 39 \times 10^{-31} \text{ kg}$					
1 dm = 0.1 m				Proton mass	m_p	$1.672\ 62 \times 10^{-27} \text{ kg}$					
1 cal (thermochemical) = 4.184 J				Neutron mass	m_n	$1.674\ 93 \times 10^{-27} \text{ kg}$					
dipole moment: 1 Debye = 3.335 64x10 ⁻³⁰ C m				Vacuum permittivity	$\epsilon_0 = \mu_0^{-1} c^{-2}$	$8.854\ 188 \times 10^{-12} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$					
force: $1N=1J \text{ m}^{-1} = 1kgms^{-2} = 10^5 \text{ dyne}$ pressure: $1Pa=1Nm^{-2} = 1Jm^{-3}$				Vacuum permeability	μ_0	$4\pi \times 10^{-7} \text{ Js}^2 \text{ C}^{-2} \text{ m}^{-1}$					
$1J = 1Nm$				Bohr magneton	$\mu_B = \frac{e\hbar}{2m_e}$	$9.274\ 02 \times 10^{-24} \text{ JT}^{-1}$					
power: $1W = 1J \text{ s}^{-1}$ potential: $1V = 1J \text{ C}^{-1}$				Nuclear magneton	$\mu_N = \frac{e\hbar}{2m_p}$	$5.05079 \times 10^{-27} \text{ JT}^{-1}$					
magnetic flux: $1T=1Vsm^{-2} = 1JCs^{-2}$ current: $1A=1Cs^{-1}$				Gravitational constant	G	$6.67259 \times 10^{-11} \text{ Nm}^2 \text{ kg}^{-2}$					
Prefixes:				Gravitational	g	9.80665 ms^{-2}					
p	n	m	m	c	d	k	M	G	acceleration		
pico	nano	micro	milli	centi	deci	kilo	mega	giga	Bohr radius	a_0	$5.291\ 77 \times 10^{-11} \text{ m}$
10 ⁻¹²	10 ⁻⁹	10 ⁻⁶	10 ⁻³	10 ⁻²	10 ⁻¹	10 ³	10 ⁶	10 ⁹			

Standard molar enthalpies of formation at 298.15 K

Temperature dependence of heat capacities, $C_{p,m} = a + bT + cT^{-2}$

M_r	$\Delta H_f^\ominus / \text{KJ/mol}$	M_r	$\Delta H_f^\ominus / \text{KJ/mol}$	$a / \text{J K}^{-1} \text{mol}^{-1}$	$b / 10^{-3} \text{J K}^{-2} \text{mol}^{-1}$	$c / 10^5 \text{J K mol}^{-1}$
H ₂ O(g)	18.015	-241.8	O ₃ (g)	47.998	+142.7	
H ₂ O(l)	18.015	-285.8	NO(g)	30.006	+90.2	
H ₂ O ₂ (l)	34.015	-187.8	NO ₂ (g)	46.006	+33.2	
NH ₃ (g)	17.031	-46.1	N ₂ O ₄ (g)	92.012	+9.2	
N ₂ H ₄ (l)	32.045	+50.6	SO ₂ (g)	64.063	-296.8	
N ₂ H(l)	43.028	+264.1	H ₂ S(g)	34.080	-20.6	
N ₂ H(g)	43.028	+294.1	SF ₆ (g)	146.054	-1209	
HNO ₃ (l)	63.013	-174.1	HF(g)	20.006	-271.1	
NH ₂ OH(s)	33.030	-114.2	HCl(g)	36.461	-92.3	
NH ₄ Cl(s)	53.492	-314.4	HCl(aq)	36.461	-167.2	
HgCl ₂ (s)	271.50	-224.3	HBr(g)	80.917	+36.4	
H ₂ SO ₄ (l)	98.078	-814.0	HI(g)	127.912	+26.5	
H ₂ SO ₄ (aq)	98.078	-909.3	CO ₂ (g)	44.010	-393.5	
NaCl(s)	58.443	-411.0	CO(g)	28.011	-110.5	
NaOH(s)	39.997	-426.7	Al ₂ O ₃ (α ,s)	101.945	-1675.7	
KCl(s)	74.555	-435.9	SiO ₂ (s)	60.085	-910.9	
KBr(s)	119.011	-392.2	FeS(s)	87.91	-100.0	
KI(s)	166.006	-327.6	FeS ₂ (s)	119.975	-178.2	
DIATOMICS	Eg. N₂, O₂, H₂	0	AgCl(s)	143.323	-127.1	
Enthalpies of fusion and evaporation $\Delta H_m / \text{KJ/mol}$ at the transition temperature				M_r	$\Delta H_f^\ominus / \text{KJ/mol}$	$\Delta H_c^\ominus / \text{KJ/mol}$
	T_f/K	Fusion^a	T_b/K	Evaporation^b		
He	3.5	0.021	4.22	0.084	16.043	-74.81
Ar	83.81	1.188	87.29	6.506	26.038	+226.8
H ₂	13.96	0.117	20.38	0.9163	28.054	+52.30
N ₂	63.15	0.719	77.35	5.586	30.070	-84.64
O ₂	54.36	0.444	90.18	6.820	42.081	53.35
Cl ₂	172.12	6.406	239.05	20.410	42.081	20.5
Br ₂	285.90	10.573	332.35	29.45	58.124	-126.11
I ₂	386.75	15.52	458.39	41.80	72.151	-146.4
Hg	234.29	2.292	629.73	59.296	84.163	-158.2
Ag	1234	11.30	2436	250.63	86.178	-198.7
Na	370.95	2.601	1156	98.01	88.115	+48.99
CO ₂	217.0	8.33	194.64	25.23 ^L	78.115	+48.99
H ₂ O	273.15	6.008	373.15	40.656 (44.016 at 298.15 K)	114.233	-249.8
NH ₃	195.40	5.652	239.73	23.351	128.175	+78.53
H ₂ S	187.61	2.377	212.80	18.673	32.042	-239.0
CH ₄	90.68	0.941	111.66	8.18	44.054	-166.0
C ₂ H ₆	89.85	2.86	184.55	14.7	46.070	-277.0
C ₆ H ₆	278.65	10.59	353.25	30.8	60.053	-484.2
CH ₃ OH	175.25	3.159	337.22	35.27 (37.99 at 298.15K)	88.107	-486.6
					88.107	-486.6
					94.114	-165.0
					93.129	-31.1
					60.056	-333.0
					75.068	-537.2
					180.159	-1274
					180.159	-1268
					342.303	-2222
					90.079	-694.0
						-1344

^aSublimation: ^avarious pressures: ^bat 1atm

Source: American Institute of Physics handbook, McGraw-Hill.

Heat capacities at 25°C

	$C_{v,m}$	$C_{p,m}$
	$\text{JK}^{-1} \text{mol}^{-1}$	$\text{JK}^{-1} \text{mol}^{-1}$
He, Ne, Ar, Kr, Xe	12.47	20.78
H ₂	20.50	28.81
O ₂	21.01	29.33
N ₂	20.83	29.14
CO ₂	28.83	37.14
NH ₃	27.17	35.48
CH ₄	27.43	35.74
N ₂ O ₄		77.28
NO ₂		37.20

F.P Depression, B.P. Elevation

Solvent	F.P °C	K _f °C kg mol ⁻¹	B.P (°C, 101kNm ⁻²)	K _b °C kg mol ⁻¹
Water	0	1.86	100.0	0.52
Benzene	5.51	5.10	80.1	2.60
Acetic Acid	16.6	3.90	118.1	3.10
Cyclohexane	6.5	20.2	81.4	2.79
Camphor	177.7	40.0	205	-
Nitrobenzene	5.7	6.9	210.9	5.24
Ethanol	-177		78.5	1.22
Chloroform	-64		61.3	3.63

Third Law entropies at 25°C, Sm⁰/J K⁻¹ mol⁻¹

Solids		Liquids		Gases	
Ag	42.68	Hg	76.02	H ₂	130.6
C(gr)	5.77	Br ₂	152.3	N ₂	192.1
C(d)	2.44			O ₂	205.1
Cu	33.4			Cl ₂	223.0
Zn	41.6	H ₂ O	70.0		
I ₂	116.7			CO ₂	213.7
S(Rh)	31.9	HNO ₃	155.6	HCl	186.8
				H ₂ S	205.6
AgCl	96.2	C ₂ H ₅ OH	161.0	NH ₃	192.5
AgBr	104.6	CH ₃ OH	126.7	CH ₄	186.1
CuSO ₄ ·5H ₂ O	305.4	C ₆ H ₆	49.03	C ₂ H ₆	229.4
HgCl ₂	144	CH ₃ COOH	159.8	CH ₃ CHO	265.7
Sucrose	360.2	C ₆ H ₁₂	298.2		

Standard molar Gibbs free energy and molar entropy of formation at 298.15 K

	M_r	$\Delta G_f^\theta/\text{KJ/mol}$	$S^\theta/\text{J K}^{-1} \text{mol}^{-1}$		M_r	$\Delta G_f^\theta/\text{KJ/mol}$	$S^\theta/\text{J K}^{-1} \text{mol}^{-1}$
H ₂ O(g)	18.015	-228.57	188.83	O ₃ (g)	47.998	163.2	238.93
H ₂ O(l)	18.015	-120.35	109.6	NO(g)	30.006	86.55	210.76
H ₂ O ₂ (l)	34.015	-120.35	109.6	NO ₂ (g)	46.006	51.31	240.06
NH ₃ (g)	17.031	-16.45	192.45	N ₂ O ₄ (g)	92.012	97.89	304.29
N ₂ H ₄ (l)	32.045	149.43	121.21	SO ₂ (g)	64.063	-300.19	248.22
N ₃ H(l)	43.028	327.3	140.6	H ₂ S(g)	34.080	-33.56	205.79
N ₃ H(g)	43.028	328.1	238.97	SF ₆ (g)	146.054	-1105.3	291.82
HNO ₃ (l)	63.013	-80.71	155.60	HF(g)	20.006	-273.2	173.78
NH ₂ OH(s)	33.030			HCl(g)	36.461	-95.30	186.91
NH ₄ Cl(s)	53.492	-202.87	94.6	HCl(aq)	36.461	-131.23	56.5
HgCl ₂ (s)	271.50	-178.6	146.0	HBr(g)	80.917	-53.45	198.70
H ₂ SO ₄ (l)	98.078	-690.00	156.90	HI(g)	127.912	1.70	206.59
H ₂ SO ₄ (aq)	98.078	-744.53	20.1	CO ₂ (g)	44.010	-394.36	213.74
NaCl(s)	58.443	-384.14	72.13	CO(g)	28.011	-137.17	197.67
NaOH(s)	39.997	-379.49	64.46	Al ₂ O ₃ (s)	101.945	-1582.3	50.92
KCl(s)	74.555	-409.14	82.59	SiO ₂	60.09	-856.64	41.84
KBr(s)	119.011	-380.66	95.90	FeS(s)	87.91	-100.4	60.29
KI(s)	166.006	-324.89	106.32	FeS ₂ (s)	119.975	-166.9	52.93
				AgCl(s)	143.323	-109.79	96.2
He(g)	4.003	0	126.15	Hg(g)	200.59	31.82	174.96
Ar(g)	39.95	0	154.84	Hg(l)	200.59	0	76.02
H ₂ (g)	2.016	0	130.684	Ag(g)	107.87	245.65	173.00
N ₂ (g)	28.013	0	191.61	Ag(s)	107.87	0	42.55
O ₂ (g)	31.999	0	205.138	Na(g)	370.95	76.76	153.71
O ₃ (g)	47.998	163.2	238.93	Na(s)	22.99	0	51.21
Cl ₂ (g)	70.91	0	223.07				
Br ₂ (g)	159.82	3.110	245.46				
Br ₂ (l)	159.82	0	152.23				
I ₂ (g)	253.81	19.33	260.69				
I ₂ (s)	253.81	0	116.135				

	M_r	$\Delta G_f^\theta/\text{KJ/mol}$	$S^\theta/\text{J K}^{-1} \text{mol}^{-1}$
organic compounds			
CH ₄ (g) methane	16.043	-50.72	186.26
C ₂ H ₂ (g) ethyne	26.038	209.20	200.94
C ₂ H ₄ (g) ethene	28.05	68.15	219.56
C ₂ H ₆ (g) ethane	30.070	-32.82	229.60
C ₃ H ₆ cyclopropane(g)	42.081	104.45	237.55
C ₃ H ₆ propene(g)	42.081	62.78	267.05
C ₄ H ₁₀ n-butane (g)	58.124	-17.03	310.23
C ₅ H ₁₂ n-pentane(g)	72.151	-8.20	348.40
C ₆ H ₁₂ cyclohexane (l)	84.163	26.8	
C ₆ H ₁₄ n-hexane (l)	86.178		204.3
C ₆ H ₆ benzene (l)	78.115	124.3	173.3
C ₆ H ₆ benzene (g)	78.115	129.72	269.31
C ₈ H ₁₈ n-octane (l)	114.233	6.4	361.1
C ₁₀ H ₈ naphthalene (l)	128.175		
CH ₃ OH (g)	32.042	-161.96	239.81
CH ₃ OH (l)	32.042	-166.27	126.8
CH ₃ CHO (g)	44.054	-128.86	250.3
CH ₃ CH ₂ OH (l)	46.07	-174.78	160.7
CH ₃ COOH (l)	60.053	-389.9	159.8
CH ₃ COOC ₂ H ₅ (l)	88.107	-332.7	259.4
C ₆ H ₅ OH (s)	94.114	-50.9	146.0
C ₆ H ₅ NH ₂ (l)	93.129		
CH ₂ (NH ₂)CO ₂ H, glycine (s)	75.068	-373.4	103.5
C ₆ H ₁₂ O ₆ , α-D-glucose (s)	180.159		
C ₆ H ₁₂ O ₆ , β-D-glucose (s)	180.159	-910	212
C ₁₂ H ₂₂ O ₁₁ , sucrose (s)	342.303	-1543	360.2
CH ₃ CH(OH)COOH	90.079		
lactic acid (s)			

Source: American Institute of Physics handbook, McGraw-Hill.