

University of Eswatini
Faculty of Science and Engineering
Department of Chemistry

Main Examination 2018/2019

Title of Paper : Applied Thermodynamics

Course code : CHE 242

Time : 3 hours

Instructions : Answer **Question 1** and any other 3 Questions
Data sheets are provided with this examination

Do not open this paper until permission has been given by the Invigilator

Question 1 [40 Marks] [Compulsory]

- a) Differentiate between the strength of an acid/base and the pH/pOH [10]
- b) An aqueous solution of KI has a freezing point of $-1.95\text{ }^{\circ}\text{C}$ and an osmotic pressure of 25.0 atm at $25.0\text{ }^{\circ}\text{C}$. Assuming that the KI completely dissociates in water, what is the density of the solution given that the freezing point constant is $1.86\text{ }^{\circ}\text{C/molal}$? [6]
- c) Derive the Clapeyron Equation showing all important steps [6]
- d) Write short notes on the following;
- (i) λ transition [3]
 - (ii) van't Hoff Factor [2]
 - (iii) Raoult's Law [3]
- e) Discuss your understanding of equilibrium for (i) chemical equilibrium, (ii) phase equilibrium and (iii) Solubility [10]

Question 2 [20 Marks]

- a) Using a rough sketch, show the important components of a phase diagram. [5]
- b) Illustrate the schematic temperature dependence of the chemical potential with temperature for the three phases of a chemical substance [5]
- c) Derive the vapor pressure of a pressurized liquid, with an aid of diagrams where necessary [5]
- d) Discuss the solubility of glucose in water [5]

Question 3 [20 Marks]

- a) Show your understanding of colligative properties by using 1 real life examples to show the use of any two scenarios of your choice. [3]
- b) Write short notes on the following;
- (i) Osmotic pressure [4]
 - (ii) Boiling point elevation [4]
 - (iii) Vapour pressure lowering [4]

- c) At 286 K, the osmotic pressure of a glucose solution is 9.97 atm. What is the freezing point depression (the density of the solution is 1.12 g/mL) given that $K_f = 1.86^\circ\text{C kg/mol}$? [5]

Question 4 [20 Marks]

- a) Derive the Gibbs energy of mixing of perfect gases [6]
- b) What is meant by the auto-ionisation of water? [5]
- c) For the equilibrium of 500 mg of vitamin C in water ($M_r = 176.826 \text{ g/mol}$) in 100 mL of water, given that the ionisation constant of vitamin C is 8.0×10^{-5} ;
- (i) Calculate the percent ionisation at equilibrium [6]
- (ii) What would be the pH of the solution at equilibrium? [3]

Question 5 [20 Marks]

- a) Using an example of your choice, demonstrate your understanding of the principle of Le Chatelier [4]
- b) Qualify the equation below; [5]

$$\Delta G = RT \ln \left(\frac{Q}{K} \right)$$

where Q is the reaction quotient and K the equilibrium constant. [5]

- c) Derive the Gibbs-Duhem equation [6]

The End

General data and fundamental constants

Quantity	Symbol	Value
Speed of light	c	$2.997\,924\,58 \times 10^8 \text{ m s}^{-1}$
Elementary charge	e	$1.602\,177 \times 10^{-19} \text{ C}$
Faraday constant	$F = N_A e$	$9.6485 \times 10^4 \text{ C mol}^{-1}$
Boltzmann constant	k	$1.380\,66 \times 10^{-23} \text{ J K}^{-1}$
Gas constant	$R = N_A k$	$8.314\,51 \text{ J K}^{-1} \text{ mol}^{-1}$
		$8.205\,78 \times 10^{-2} \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$
		$6.2364 \times 10 \text{ L Torr K}^{-1} \text{ mol}^{-1}$
Planck constant	h	$6.626\,08 \times 10^{-34} \text{ J s}$
	$\hbar = h/2\pi$	$1.054\,57 \times 10^{-34} \text{ J s}$
Avogadro constant	N_A	$6.022\,14 \times 10^{23} \text{ mol}^{-1}$
Atomic mass unit	u	$1.660\,54 \times 10^{-27} \text{ Kg}$
Mass		
electron	m_e	$9.109\,39 \times 10^{-31} \text{ Kg}$
proton	m_p	$1.672\,62 \times 10^{-27} \text{ Kg}$
neutron	m_n	$1.674\,93 \times 10^{-27} \text{ Kg}$
Vacuum permittivity	$\epsilon_0 = 1/c^2 \mu_0$	$8.854\,19 \times 10^{-12} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$
	$4\pi\epsilon_0$	$1.112\,65 \times 10^{-10} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$
Vacuum permeability	μ_0	$4\pi \times 10^{-7} \text{ J s}^2 \text{ C}^{-2} \text{ m}^{-1}$
		$4\pi \times 10^{-7} \text{ T}^2 \text{ J}^{-1} \text{ m}^3$
Magneton		
Bohr	$\mu_B = e\hbar/2m_e$	$9.274\,02 \times 10^{-24} \text{ J T}^{-1}$
nuclear	$\mu_N = e\hbar/2m_p$	$5.050\,79 \times 10^{-27} \text{ J T}^{-1}$
g value	g_e	2.002 32
Bohr radius	$a_0 = 4\pi\epsilon_0 \hbar^2 / m_e e^2$	$5.291\,77 \times 10^{-11} \text{ m}$
Fine-structure constant	$\alpha = \mu_0 e^2 c / 2\hbar$	$7.297\,35 \times 10^{-3}$
Rydberg constant	$R_\infty = m_e e^4 / 8\hbar^3 c \epsilon_0^2$	$1.097\,37 \times 10^7 \text{ m}^{-1}$
Standard acceleration of free fall	g	$9.806\,65 \text{ m s}^{-2}$
Gravitational constant	G	$6.672\,59 \times 10^{-11} \text{ N m}^2 \text{ Kg}^{-2}$

Conversion factors

1 cal =	4.184 joules (J)	1 erg =	$1 \times 10^{-7} \text{ J}$
1 eV =	$1.602\,2 \times 10^{-19} \text{ J}$	1 eV/molecule =	$96\,485 \text{ kJ mol}^{-1}$

Prefixes	f	p	n	μ	m	c	d	k	M	G
	femto	pico	nano	micro	milli	centi	deci	kilo	mega	giga
	10^{-15}	10^{-12}	10^{-9}	10^{-6}	10^{-3}	10^{-2}	10^{-1}	10^3	10^6	10^9

PERIODIC TABLE OF ELEMENTS

GROUPS

PERIODS	GROUPS																	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
	IA	IIA	IIIB	IVB	VB	VIB	VIIIB		VIIIB		IB	IIIB	IIIA	IVA	VA	VIA	VIIA	VIIIA
1	1,008 H																	
2	6.941 Li 3	9.012 Be 4																
3	22.990 Na 11	24.305 Mg 12																
4	39.098 K 19	40.078 Ca 20	44.956 Sc 21	47.88 Ti 22	50.942 V 23	51.996 Cr 24	54.938 Mn 25	55.847 Fe 26	58.933 Co 27	58.69 Ni 28	63.546 Cu 29	65.39 Zn 30	69.723 Ga 31	72.61 Ge 32	74.922 As 33	78.96 Se 34	79.904 Br 35	83.80 Kr 36
5	85.468 Rb 37	87.62 Sr 38	88.906 Y 39	91.224 Zr 40	92.906 Nb 41	95.94 Mo 42	98.907 Tc 43	101.07 Ru 44	102.91 Rh 45	106.42 Pd 46	107.87 Ag 47	112.41 Cd 48	114.82 In 49	118.71 Sn 50	121.75 Sb 51	127.60 Te 52	126.90 I 53	131.29 Xe 54
6	132.91 Cs 55	137.33 Ba 56	138.91 *La 57	178.49 Hf 72	180.95 Ta 73	183.85 W 74	186.21 Re 75	190.2 Os 76	192.22 Ir 77	195.08 Pt 78	196.97 Au 79	200.59 Hg 80	204.38 Tl 81	207.2 Pb 82	208.98 Bi 83	(209) Po 84	(210) At 85	(222) Rn 86
7	223 Fr 87	226.03 Ra 88	(227) **Ac 89															

Atomic mass →
Symbol →
Atomic No. →

TRANSITION ELEMENTS

140.12 Ce 58	140.91 Pr 59	144.24 Nd 60	150.36 Sm 62	(145) Pm 61	151.96 Eu 63	157.25 Gd 64	158.93 Tb 65	162.50 Dy 66	164.93 Ho 67	167.26 Er 68	168.93 Tm 69	173.04 Yb 70	174.97 Lu 71
232.04 Th 90	231.04 Pa 91	238.03 U 92	(244) Pu 94	(243) Am 95	(247) Cm 96	(247) Bk 97	(251) Cf 98	(252) Es 99	(257) Fm 100	(258) Md 101	(259) No 102	(260) Lr 103	

*Lanthanide Series

**Actinide Series

() indicates the mass number of the isotope with the longest half-life.