

UNIVERSITY OF ESWATINI

FIRST SEMESTER MAIN EXAMINATION PAPER, NOVEMBER 2019

FACULTY OF SCIENCE AND ENGINEERING

DEPARTMENT OF COMPUTER SCIENCE

COURSE CODE: CSC 205

TITLE OF PAPER: PROBABILITY AND STATISTICS

TIME ALLOWED: 3 HOURS

Instructions

1. Show all your working.
2. Answer any FIVE questions.

Special Requirements

Scientific calculator

Additional Material (s)

1. Statistical Tables

*Candidates may complete the front cover of their answer book when instructed by the Chief Invigilator and sign their examination attendance cards but must **NOT** write anything else until the start of the examination period is announced.*

No electronic devices capable of storing and retrieving text, including electronic dictionaries and any form of foreign material may be used while in the examination room.

DO NOT turn examination paper over until instructed to do so.

Question 1 [6+8+4+2 marks]

A study of the effects of smoking on sleep patterns is conducted. The measure observed is the time, in minutes, that it takes to fall asleep. These data are obtained:

Smokers: 69.3 56.0 22.1 47.6
 53.2 48.1 52.7 34.4
 60.2 43.8 23.2 13.8

Nonsmokers: 28.6 25.1 26.4 34.9
 29.8 28.4 38.5 30.2
 30.6 31.8 41.6 21.1
 36.0 37.9 13.9

- Find the sample mean for each group.
- Find the sample standard deviation for each group.
- Make a dot plot of the data sets A (Smokers) and B (Nonsmokers) on the same line.
- Comment on what kind of impact smoking appears to have on the time required to fall asleep.

Question 2 [2+2+3+2+5+3+3 marks]

- State the additive law of probability.
- Suppose the manufacturer's specifications for the length of a certain type of computer cable are 2000 ± 10 millimeters. In this industry, it is known that small cable is just as likely to be defective (not meeting specifications) as large cable. That is, the probability of randomly producing a cable with length exceeding 2010 millimeters is equal to the probability of producing a cable with length smaller than 1990 millimeters. The probability that the production procedure meets specifications is known to be 0.99.
 - What is the probability that a cable selected randomly is too large?
 - What is the probability that a randomly selected cable is larger than 1990 millimeters?
- State (without proof) the law of total probability.
- State and prove the Bayes rule
- In a certain assembly plant, three machines, B_1 , B_2 , and B_3 , make 30%, 45%, and 25%, respectively, of the products. It is known from past experience that 2%, 3%, and 2% of the products made by each machine, respectively, are defective. Now, suppose that a finished product is randomly selected. What is the probability that it is defective?
- With reference to (e), if a product was chosen randomly and found to be defective, what is the probability

that it was made by machine B_3 ?

Question 3 [3+3+2+2+3+3+4 marks]

- a. The probability that a patient recovers from a rare blood disease is 0.4. If 15 people are known to have contracted this disease, what is the probability that (i) at least 10 survive, (ii) from 3 to 8 survive, and (iii) exactly 5 survive?
- b. Let Y be a random variable with $p(y)$ given in the table below. Find $E(Y)$, $E(1/Y)$, $E(Y^2 - 1)$ and $V(Y)$.

y	1	2	3	4
$p(y)$	0.4	0.3	0.2	0.1

Question 4 [3+3+4+3+4+3 marks]

- a. Suppose that Y has density function

$$f(y) = \begin{cases} ky(1-y), & 0 \leq y \leq 1 \\ 0, & \text{elsewhere.} \end{cases}$$

- Find the value of k that makes $f(y)$ a probability density function.
 - Find $P(0.4 \leq Y \leq 1)$
 - Find $P(Y \leq 0.4|Y \leq 0.4)$
- b. The loaves of rye bread distributed to local stores by a certain bakery have an average length of 30 centimeters and a standard deviation of 2 centimeters. Assuming that the lengths are normally distributed, what percentage of the loaves are
- longer than 31.7 centimeters?
 - between 29.3 and 33.5 centimeters in length?
 - shorter than 25.5 centimeters?

Question 5 [5 + 12 + 3 marks]

The grades of a class of 9 students on a midterm report (x) and on the final examination (y) are as follows:

x : 77 50 71 72 81 94 96 99 67

y : 82 66 78 34 47 85 99 99 68

- Construct a scatter diagram for these data.
- Estimate the linear regression line, include the regression line on the scatter diagram in (a).

- c. Estimate the final examination grade of a student who received a grade of 85 on the midterm report.

Question 6 [4 + 10 + 6 marks]

On a factory production line it is important that the time taken to assemble a component is within certain limits. Assembly times (in minutes) for a single component are recorded for a random sample of twelve factory workers and the values are as follows.

11.7 12.8 9.9 10.6 11.6 10.6 13.1 11.2 11.6 11.9 10.9 12.7

- (a) Calculate the mean and standard deviation of these observed times.
- (b) Assuming that the underlying distribution of assembly times is Normal, calculate 95% confidence intervals for the mean and for the standard deviation of the assembly times.
- (c) In order to facilitate smooth operation of the entire production process, the assembly times must satisfy certain conditions. Test at the 5% significance level the hypothesis that the mean assembly time in the factory is 11 minutes against the hypothesis that it is greater than 11 minutes.

APPENDIX 3

Tables

Table 1 Binomial Probabilities

Tabulated values are $P(Y \leq a) = \sum_{y=0}^a p(y)$. (Computations are rounded at third decimal place.)

(a) $n = 5$

a	p													a
	0.01	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95	0.99	
0	.951	.774	.590	.328	.168	.078	.031	.010	.002	.000	.000	.000	.000	0
1	.999	.977	.919	.737	.528	.337	.188	.087	.031	.007	.000	.000	.000	1
2	1.000	.999	.991	.942	.837	.683	.500	.317	.163	.058	.009	.001	.000	2
3	1.000	1.000	1.000	.993	.969	.913	.812	.663	.472	.263	.081	.023	.001	3
4	1.000	1.000	1.000	1.000	.998	.990	.969	.922	.832	.672	.410	.226	.049	4

(b) $n = 10$

a	p													a
	0.01	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95	0.99	
0	.904	.599	.349	.107	.028	.006	.001	.000	.000	.000	.000	.000	.000	0
1	.996	.914	.736	.376	.149	.046	.011	.002	.000	.000	.000	.000	.000	1
2	1.000	.988	.930	.678	.383	.167	.055	.012	.002	.000	.000	.000	.000	2
3	1.000	.999	.987	.879	.650	.382	.172	.055	.011	.001	.000	.000	.000	3
4	1.000	1.000	.998	.967	.850	.633	.377	.166	.047	.006	.000	.000	.000	4
5	1.000	1.000	1.000	.994	.953	.834	.623	.367	.150	.033	.002	.000	.000	5
6	1.000	1.000	1.000	.999	.989	.945	.828	.618	.350	.121	.013	.001	.000	6
7	1.000	1.000	1.000	1.000	.998	.988	.945	.833	.617	.322	.070	.012	.000	7
8	1.000	1.000	1.000	1.000	1.000	.998	.989	.954	.851	.624	.264	.086	.004	8
9	1.000	1.000	1.000	1.000	1.000	1.000	.999	.994	.972	.893	.651	.401	.096	9


Table 1 (Continued)
(c) $n = 15$

a	p												a	
	0.01	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95		0.99
0	.860	.463	.206	.035	.005	.000	.000	.000	.000	.000	.000	.000	.000	0
1	.990	.829	.549	.167	.035	.005	.000	.000	.000	.000	.000	.000	.000	1
2	1.000	.964	.816	.398	.127	.027	.004	.000	.000	.000	.000	.000	.000	2
3	1.000	.995	.944	.648	.297	.091	.018	.002	.000	.000	.000	.000	.000	3
4	1.000	.999	.987	.836	.515	.217	.059	.009	.001	.000	.000	.000	.000	4
5	1.000	1.000	.998	.939	.722	.403	.151	.034	.004	.000	.000	.000	.000	5
6	1.000	1.000	1.000	.982	.869	.610	.304	.095	.015	.001	.000	.000	.000	6
7	1.000	1.000	1.000	.996	.950	.787	.500	.213	.050	.004	.000	.000	.000	7
8	1.000	1.000	1.000	.999	.985	.905	.696	.390	.131	.018	.000	.000	.000	8
9	1.000	1.000	1.000	1.000	.996	.966	.849	.597	.278	.061	.002	.000	.000	9
10	1.000	1.000	1.000	1.000	.999	.991	.941	.783	.485	.164	.013	.001	.000	10
11	1.000	1.000	1.000	1.000	1.000	.998	.982	.909	.703	.352	.056	.005	.000	11
12	1.000	1.000	1.000	1.000	1.000	1.000	.996	.973	.873	.602	.184	.036	.000	12
13	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.995	.965	.833	.451	.171	.010	13
14	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.995	.965	.794	.537	.140	14

(d) $n = 20$

a	p												a	
	0.01	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95		0.99
0	.818	.358	.122	.012	.001	.000	.000	.000	.000	.000	.000	.000	.000	0
1	.983	.736	.392	.069	.008	.001	.000	.000	.000	.000	.000	.000	.000	1
2	.999	.925	.677	.206	.035	.004	.000	.000	.000	.000	.000	.000	.000	2
3	1.000	.984	.867	.411	.107	.016	.001	.000	.000	.000	.000	.000	.000	3
4	1.000	.997	.957	.630	.238	.051	.006	.000	.000	.000	.000	.000	.000	4
5	1.000	1.000	.989	.804	.416	.126	.021	.002	.000	.000	.000	.000	.000	5
6	1.000	1.000	.998	.913	.608	.250	.058	.006	.000	.000	.000	.000	.000	6
7	1.000	1.000	1.000	.968	.772	.416	.132	.021	.001	.000	.000	.000	.000	7
8	1.000	1.000	1.000	.990	.887	.596	.252	.057	.005	.000	.000	.000	.000	8
9	1.000	1.000	1.000	.997	.952	.755	.412	.128	.017	.001	.000	.000	.000	9
10	1.000	1.000	1.000	.999	.983	.872	.588	.245	.048	.003	.000	.000	.000	10
11	1.000	1.000	1.000	1.000	.995	.943	.748	.404	.113	.010	.000	.000	.000	11
12	1.000	1.000	1.000	1.000	.999	.979	.868	.584	.228	.032	.000	.000	.000	12
13	1.000	1.000	1.000	1.000	1.000	.994	.942	.750	.392	.087	.002	.000	.000	13
14	1.000	1.000	1.000	1.000	1.000	.998	.979	.874	.584	.196	.011	.000	.000	14
15	1.000	1.000	1.000	1.000	1.000	1.000	.994	.949	.762	.370	.043	.003	.000	15
16	1.000	1.000	1.000	1.000	1.000	1.000	.999	.984	.893	.589	.133	.016	.000	16
17	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.996	.965	.794	.323	.075	.001	17
18	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.999	.992	.931	.608	.264	.017	18
19	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.999	.988	.878	.642	.182	19


Table 4 Normal Curve Areas
 Standard normal probability in right-hand tail
 (for negative values of z , areas are found by symmetry)



z	Second decimal place of z									
	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
2.9	.0019	.0018	.0017	.0017	.0016	.0016	.0015	.0015	.0014	.0014
3.0	.00135									
3.5	.000233									
4.0	.0000317									
4.5	.00000340									
5.0	.000000287									

From R. E. Walpole, *Introduction to Statistics* (New York: Macmillan, 1968).

Table 5 Percentage Points of the *t* Distributions



$t_{.100}$	$t_{.050}$	$t_{.025}$	$t_{.010}$	$t_{.005}$	df
3.078	6.314	12.706	31.821	63.657	1
1.886	2.920	4.303	6.965	9.925	2
1.638	2.353	3.182	4.541	5.841	3
1.533	2.132	2.776	3.747	4.604	4
1.476	2.015	2.571	3.365	4.032	5
1.440	1.943	2.447	3.143	3.707	6
1.415	1.895	2.365	2.998	3.499	7
1.397	1.860	2.306	2.896	3.355	8
1.383	1.833	2.262	2.821	3.250	9
1.372	1.812	2.228	2.764	3.169	10
1.363	1.796	2.201	2.718	3.106	11
1.356	1.782	2.179	2.681	3.055	12
1.350	1.771	2.160	2.650	3.012	13
1.345	1.761	2.145	2.624	2.977	14
1.341	1.753	2.131	2.602	2.947	15
1.337	1.746	2.120	2.583	2.921	16
1.333	1.740	2.110	2.567	2.898	17
1.330	1.734	2.101	2.552	2.878	18
1.328	1.729	2.093	2.539	2.861	19
1.325	1.725	2.086	2.528	2.845	20
1.323	1.721	2.080	2.518	2.831	21
1.321	1.717	2.074	2.508	2.819	22
1.319	1.714	2.069	2.500	2.807	23
1.318	1.711	2.064	2.492	2.797	24
1.316	1.708	2.060	2.485	2.787	25
1.315	1.706	2.056	2.479	2.779	26
1.314	1.703	2.052	2.473	2.771	27
1.313	1.701	2.048	2.467	2.763	28
1.311	1.699	2.045	2.462	2.756	29
1.282	1.645	1.960	2.326	2.576	inf.

From "Table of Percentage Points of the *t*-Distribution." Computed by Maxine Merrington, *Biometrika*, Vol. 32 (1941), p. 300.