

UNIVERSITY OF SWAZILAND

SUPPLEMENTARY EXAMINATIONS 2006

BSc. / BEd. / B.A.S.S. IV

TITLE OF PAPER : FLUID DYNAMICS

COURSE NUMBER : M 455

TIME ALLOWED : THREE (3) HOURS

INSTRUCTIONS : 1. THIS PAPER CONSISTS OF
SEVEN QUESTIONS.
2. ANSWER ANY FIVE QUESTIONS

SPECIAL REQUIREMENTS : NONE

THIS EXAMINATION PAPER SHOULD NOT BE OPENED UNTIL
PERMISSION HAS BEEN GRANTED BY THE INVIGILATOR.

QUESTION 1

1. (a) Show that the velocity field for a fluid motion given by

$$\mathbf{q} = (x + 2y + 4z)\mathbf{i} + (2x - 3y - z)\mathbf{j} + (4x - y + 2z)\mathbf{k}$$

represents a possible motion of an incompressible and irrotational fluid.

[5 marks]

- (b) Find the velocity potential for the flow in (a).

[5 marks]

- (c) Consider the two-dimensional velocity field given by

$$\mathbf{q} = \frac{y}{x^2 - 1}\mathbf{i} - \frac{x}{x^2 - 1}\mathbf{j}$$

Determine the equation of the streamline passing through the point (4, 3)

[5 marks]

- (d) Find the velocity components for a flow with velocity potential given by

$$\phi = \frac{x}{x^2 + y^2}$$

[5 marks]

QUESTION 2

2. Consider the boundary layer equations in the form

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} = 0 \quad (1)$$

$$u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} = \nu \frac{\partial^2 u}{\partial y^2} - au \quad (2)$$

with boundary conditions

$$u = ax, \quad v = -(\nu a)^{\frac{1}{2}} \quad \text{on } y = 0 \quad \text{and} \quad u = 0 \quad \text{on } y = \infty$$

Using the similarity transformation $\eta = y \left(\frac{a}{\nu}\right)^{\frac{1}{2}}$ and the stream function formulation $\psi = -x(\nu a)^{\frac{1}{2}} f(\eta)$ where a is a constant and ν is the dynamic viscosity, Show that equation (2) and the boundary conditions can be transformed into

$$f''' + ff'' - (f')^2 - f' = 0$$

$$f = 1, \quad f' = 1 \quad \text{on } \eta = 0 \quad \text{and} \quad f' = 0 \quad \text{on } \eta = \infty$$

[20 marks]

QUESTION 3

3. A viscous liquid occupies the space between two coaxial, infinitely-long cylinders. The inner cylinder has radius a and is fixed, while the outer cylinder has radius b and is rotating with constant angular velocity Ω . Let (r, θ, z) be cylindrical polar coordinates with z -axis coinciding with the cylinders' axis such that the outer cylinder is rotating in the direction of increasing θ . Assuming that the velocity of the liquid has the form $\mathbf{q} = u(r)\hat{\theta}$ (where $u(r)$ means that u is a function of r only and $\hat{\theta}$ is a unit vector in the θ direction) and that body forces are negligible, use the Navier-Stokes equations in the form

$$\nabla \left(\frac{1}{2} \mathbf{q}^2 \right) - \mathbf{q} \times (\nabla \times \mathbf{q}) = -\frac{1}{\rho} \nabla p - \nu \nabla \times (\nabla \times \mathbf{q})$$

to show that $u = A(r - a^2/r)$, where $A = \Omega(1 - a^2/b^2)$.

[20 marks]

QUESTION 4

4. (a) The velocity potential for a steady incompressible, irrotational flow with circulation around a fixed cylinder of radius a is given in cylindrical polar coordinates by

$$\phi = Ur \left(1 + \frac{a^2}{r^2} \right) \cos \theta + \frac{k\theta}{2\pi}$$

where U is the uniform speed at infinity. Find the corresponding

(i) velocity field \mathbf{q} . [4 marks]

(ii) stream function ψ . [6 marks]

- (b) In the two $z = x + iy$ plane, a line vortex of strength $m > 0$, is placed at $z = c$ and another, of strength $-m$, at $z = -c$, where c is a real positive number. Both vortices are held fixed at these locations. Write down the complex potential w for this flow and show that the stream function ψ and the velocity potential ϕ are given by

$$\psi = \frac{m}{4\pi} \log \frac{(x-c)^2 + y^2}{(x+c)^2 + y^2} \quad \text{and} \quad \phi = -\frac{m}{2\pi} \tan^{-1} \frac{2cy}{x^2 + y^2 - c^2}$$

hint: You may set $z - c = r_1 e^{i\theta_1}$ and $z + c = r_2 e^{i\theta_2}$, and

use $\tan^{-1} A + \tan^{-1} B = \tan^{-1} \left(\frac{A+B}{1-AB} \right)$ [10 marks]

QUESTION 5

5. (a) Show that the complex velocity potential corresponding to the velocity field

$$\mathbf{q} = 3(y^2 - x^2)\mathbf{i} + 6xy\mathbf{j} \text{ is } w(z) = z^3$$

[6 marks]

- (b) A fluid has the complex velocity potential

$$w(z) = i + z^2, \quad z = x + iy$$

Find the stream function and the velocity potential for this flow [4 marks]

- (c) Consider the viscous flow of fluid confined between two parallel flat plates of infinite extent in the xy plane. The distance between the plates is h with the lower plate fixed at $y = -h/2$ and the upper plate is fixed at $y = h/2$. If the velocity field for the flow is of the form

$$\mathbf{q} = (u(y), 0, 0)$$

use Navier-Stokes equations in the form

$$\frac{\partial \mathbf{q}}{\partial t} + (\mathbf{q} \cdot \nabla) \mathbf{q} = -\frac{1}{\rho} \nabla p + \nu \nabla^2 \mathbf{q}$$

to show that the velocity profile for this flow is

$$u(y) = \frac{1}{2\mu} \frac{dp}{dx} \left\{ y^2 - \frac{h^2}{4} \right\}$$

[10 marks]

QUESTION 6

6. (a) An incompressible fluid flows steadily past a sphere of radius a located at the origin. The velocity at large distances away from the sphere is $U\mathbf{i}$, where \mathbf{i} is a unit vector in the x -direction. It is assumed that the motion is irrotational and that there are no body forces. Given that the velocity potential for the flow is given by

$$\phi = U \left(r + \frac{a^3}{2r^2} \right) \cos \theta$$

in the usual spherical coordinates (r, θ, ψ) , with $\theta = 0$, measured from the x -axis.

- i. Find the velocity components of the flow. [5 marks]
ii. Show that the pressure is given by

$$p = p_\infty + \frac{1}{2}\rho U^2 \left(1 - \frac{9}{4} \sin^2 \theta \right)$$

where p_∞ is the pressure at large distances from the sphere. [5 marks]

- (b) At a point in a steady, incompressible fluid having cylindrical coordinates (r, θ, z) the velocity components are

$$(r^2 \cos \theta, -3r^2 \sin \theta, 0)$$

Determine whether or not the equation of continuity is satisfied, and if so find the equations of the streamlines [10 marks]

QUESTION 7

7. (a) The velocity field for an inviscid fluid is given by

$$u = -ay, \quad v = ax, \quad w = 0$$

where a is a constant. Assuming that there are no body forces acting on the fluid,

- i. Prove that the flow is incompressible. [2 marks]
- ii. Find the rotation of the fluid. [3 marks]
- iii. If the pressure at $x = y = 0$ is p_0 , prove that the pressure at each point of the fluid is given by

$$p = p_0 + \frac{1}{2}\rho a^2(x^2 + y^2)$$

[7 marks]

- (b) Find the complex velocity potential for a fluid moving with constant speed V_0 in a direction making an angle α with the positive x -axis. [4 marks]
- (c) Determine the equations for the velocity potential ϕ , stream function ψ for the flow in (b). [4 marks]