

UNIVERSITY OF SWAZILAND

FACULTY OF SCIENCE

DEPARTMENT OF PHYSICS

MAIN EXAMINATION 2008/2009

TITLE OF PAPER : ELECTRICITY & MAGNETISM

COURSE NUMBER : P221

TIME ALLOWED : THREE HOURS

**INSTRUCTIONS : ANSWER ANY FOUR OUT OF
SIX QUESTIONS**

**EACH QUESTION CARRIES 25
MARKS**


**MARKS FOR DIFFERENT
SECTIONS OF EACH QUESTION
ARE SHOWN IN THE
RIGHT-HAND MARGIN**

THIS PAPER HAS 8 PAGES, INCLUDING THIS PAGE

**DO NOT OPEN THE PAPER UNTIL PERMISSION HAS BEEN
GIVEN BY THE INVIGILATOR**

Question 1

- a) Describe the physical content of Kirchoff's current law and Kirchoff's voltage law. [6]



- b) In the circuit drawn above, the battery provides a steady voltage of 10 volts. The values of the resistances are given in the figure in Ohms.

What is the current that flows through each resistance? [6]

What is the potential drop across each resistance? [6]


- c) A battery has an internal resistance. Describe an experiment that would enable you to

a) evaluate the magnitude of the internal resistance

b) determine the *emf* of the battery [7]

Question 2

- a) Consider a positive point charge of magnitude Q_1 . What is the magnitude and direction of the electric field produced by this charge at a distance r from the charge expressed in S.I. units? [2]
- b) What is the force that acts between two point charges of magnitudes Q_1 and Q_2 ? [2]
- c) What is the principle of superposition? [3]
- d) From the figure, determine the magnitude of the resultant force acting on Q_3 produced by charges Q_1 and Q_2 . The three point charges are represented by black circles. d_{32} is the distance between point charges Q_3 and Q_2 , and d_{31} is the distance between charges Q_3 and Q_1 . The line connecting Q_1 and Q_3 is inclined at 45° to the horizontal and vertical axes. [12]
- e) What angle does the resultant force make with the horizontal axis? [6]



Question 3

Write down an expression for the Lorentz force that acts on a point charge Q which is moving in a region of space where both the electric field \mathbf{E} and magnetic flux density \mathbf{B} are non-zero. [3]

- a) Describe how a charged particle with a well-defined velocity can be selected from a beam of similar particles which have a range of velocities. [12]
- b) A proton is in a uniform magnetic field of 0.1 tesla , which is perpendicular to the velocity of the proton. The proton moves in a circular orbit of radius 0.25 m . Calculate the orbital speed of the proton. [5]
- c) If the proton were to be replaced by an electron moving with the same velocity as the proton, what would the radius of the orbit of the electron be? [5]

Question 4

- a) Write down an expression for Ampère's circuital law, and hence determine the magnitude of the magnetic field H at a distance a from a very long, straight wire carrying current I . The wire is parallel to the z -direction. [6]
- b) The conducting wire carries a conventional current that flows in the positive z -direction. Draw a diagram showing the long, straight wire and the direction and orientation, with respect to the wire, of lines of H . [6]
- c) 2 long, straight wires, both parallel to the z -direction, carry a conventional current in the positive z -direction. Is there a force between them, and, if it exists, is it attractive or repulsive? [6]
- d) The direction of the conventional current is reversed in one of the wires, but its magnitude is kept the same. Explain all the differences that this change in direction of current causes. [7]

Question 5

a) A circuit contains a resistor, an inductor and a capacitor connected in series to an alternating voltage supply. What is the impedance of this circuit? Explain carefully the meaning of each of the symbols you have used in your expression for the impedance. [7]

b) For a certain angular frequency, the current is in phase with the alternating voltage. What is this frequency called: what is its magnitude, and why are the voltage and current in phase? [12]

c) The quality factor, $Q = \sqrt{\frac{L}{C}} \frac{1}{R}$. Explain the importance of Q when describing the response of the circuit to an alternating driving voltage. [6]

Question 6

What is a dielectric? What is the magnitude of the electrical conductivity of an ideal dielectric? [2]

A parallel-plate capacitor was constructed from two rectangular plates of aluminium, each of area A , separated by a distance d . A battery that provided $50V$ was used to charge the capacitor. After the capacitor was fully charged, the battery was disconnected from the capacitor, and a dielectric was inserted between the two aluminium plates, which completely filled the space between the aluminium plates. What changes were caused by the insertion of the dielectric? [6]

The dimensions of the capacitor and dielectric were as follows:

$$A = 0.02 \text{ m}^2$$

$$d = 0.025 \text{ m}$$

$$\text{thickness of dielectric} = 0.025 \text{ m}$$

$$\text{relative permittivity of the dielectric} = 4$$

Calculate the following quantities

- (a) the capacitance before the dielectric slab was inserted [5]
- (b) the charge on the plates of the capacitor before the dielectric was inserted [4]
- (c) the charge on the plates of the capacitor after the dielectric was inserted [2]
- (d) the potential difference between the plates, and the capacitance, when the dielectric was between the plates. [6]

PHYSICAL CONSTANTS AND UNITS

Acceleration due to gravity	g	9.81 m s^{-2}
Gravitational constant	G	$6.673 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
Avogadro constant	N_A	$6.022 \times 10^{23} \text{ mol}^{-1}$
(Note: 1 mole = 1 gram molecular-weight)		
Ice point	T_{ice}	273.15 K
Gas constant	R	$8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
Boltzmann constant	k, k_B	$1.381 \times 10^{-23} \text{ J K}^{-1} = 0.862 \times 10^{-4} \text{ eV K}^{-1}$
Stefan constant	σ	$5.670 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$
Rydberg constant	R_∞ $R_\infty hc$	$1.097 \times 10^7 \text{ m}^{-1}$ 13.606 eV
Planck constant	h	$6.626 \times 10^{-34} \text{ J s} = 4.136 \times 10^{-15} \text{ eV s}$
$h/2\pi$	\hbar	$1.055 \times 10^{-34} \text{ J s} = 6.582 \times 10^{-16} \text{ eV s}$
Speed of light in vacuo	c	$2.998 \times 10^8 \text{ m s}^{-1}$
	$\hbar c$	197.3 MeV fm
Charge of proton	e	$1.602 \times 10^{-19} \text{ C}$
Mass of electron	m_e	$9.109 \times 10^{-31} \text{ kg}$
Rest energy of electron		0.511 MeV
Mass of proton	m_p	$1.673 \times 10^{-27} \text{ kg}$
Rest energy of proton		938.3 MeV
One atomic mass unit	u	$1.66 \times 10^{-27} \text{ kg}$
Atomic mass unit energy equivalent		931.5 MeV
Electric constant	ϵ_0	$8.854 \times 10^{-12} \text{ F m}^{-1}$
Magnetic constant	μ_0	$4\pi \times 10^{-7} \text{ H m}^{-1}$
Bohr magneton	μ_B	$9.274 \times 10^{-24} \text{ A m}^2 (\text{J T}^{-1})$
Nuclear magneton	μ_N	$5.051 \times 10^{-27} \text{ A m}^2 (\text{J T}^{-1})$
Fine-structure constant	$\alpha = e^2/4\pi\epsilon_0\hbar c$	$7.297 \times 10^{-3} = 1/137.0$
Compton wavelength of electron	$\lambda_c = h/mc$	$2.426 \times 10^{-12} \text{ m}$
Bohr radius	a_0	$5.2918 \times 10^{-11} \text{ m}$
angstrom	\AA	10^{-10} m
torr (mm Hg, 0°C)	torr	133.32 Pa (N m^{-2})
barn	b	10^{-28} m^2