

UNIVERSITY OF SWAZILAND

SUPPLEMENTARY EXAMINATION PAPER 2006

TITLE OF PAPER : DESCRIPTIVE/INFERENTIAL STATISTICS
COURSE CODE : ST230/IDE-ST230-1&2
TIME ALLOWED : 3 (THREE) HOURS
**REQUIRMENTS : STATISTICAL TABLES
AND CALCULATOR**
**INSTRUCTIONS : ANSWER ANY 5 (FIVE) QUESTIONS.
ALL QUESTIONS CARRY EQUAL MARKS.**

**THIS PAPER IS NOT TO BE OPENED UNTIL PERMISSION HAS BEEN
GRANTED BY THE INVIGILATOR**

ANSWER ANY FIVE QUESTIONS:

(You must show all your works in order to obtain full marks.)

QUESTION ONE.

[2 + 4 + 8 + 3 + 3 marks]

1.1 A committee was formed with three members. Find the probability

- a. that the committee members are all men.
- b. that the committee members are either all men or all women.

1.2 A company manufactures steel rods of 500 cm length, each one having two flaws on average. The process ultimately divides these rods into 100 cm segments. What is the probability that a particular 100 cm segment will have no flaws?

1.3 In a sample of 50 people, 21 had type O blood, 22 had type A blood, 5 had B blood and 2 had type AB blood. Find the probability

- a. that a person has type A or type B blood.
- b. that a person has neither type B nor type O blood.

QUESTION TWO.

[12 + 3 + 3 + 2 marks]

2.1 Consider the following table (use 2004 as base year):

Commodity	2004		2005	
	Prices (E per kg)	Quantity Purchased	Prices (E per kg)	Quantity Purchased
A	15	10	18	15
B	5	8	6	10
C	10	5	14	8

Compute Fisher's volume index.

2.2 The following data represents the number of students were in the classroom before the lecturer arrived in a Statistics Course during a semester:

43 54 67 53 61 71 46 61 55 56
 58 60 66 54 68 52 36 64 51 52
 57 52 63 59 69 70 42 66 58 53

- a. Construct a Stem and Leaf diagram for the above data.
- b. Find the median and the mode of the above data.

QUESTION THREE.

[5 + 8 + 5 + 2 marks]

3.1 The following table shows the quarterly sales of a company in thousands of tons for a period of three years:

Sales:		Quarter			
		1	2	3	4
Year	1	32	48	54	28
	2	34	57	62	33
	3	39	62	74	39

- Compute the trend of the above sales data.
 - Find the average seasonal variation.
- 3.2 Draw a Histogram and a Frequency Polygon using the following grouped frequency distribution (consider equal classes). You must draw on your answer script; do not use any graph paper.

Class limits	Frequency
Less than 5	7
Less than 8	22
Less than 11	44
Less than 14	58
Less than 17	60

QUESTION FOUR.

[10 + 10 marks]

4.1 A researcher wishes to determine whether there is a relationship between the gender of an individual and the amount of alcohol consumed. A sample of 68 people is selected, and the following data are obtained:

Gender	Alcohol Consumption		
	Low	Moderate	High
Male	10	9	8
Female	13	16	12

At 5% level of significance, can the researcher conclude that alcohol consumption is related to gender?

4.2 A machine is supposed to be adjusted to produce components to a dimension of 2.0 cm. In a sample of 50 components, the mean was found to be 2.001 cm and the standard deviation to be 0.003 cm. Is there evidence to suggest that the machine is set to high? Use $\alpha = 0.05$.

QUESTION FIVE.

[10 + 4 + 6 marks]

Consider the following results:

$$\Sigma X = 67 \quad \Sigma Y = 225 \quad \Sigma X^2 = 645 \quad \Sigma XY = 1300 \quad \Sigma Y^2 = 5331 \quad n = 10$$

- Estimate the linear regression line.
- What would be the value of Y when X is 10.
- Compute the coefficient of correlation (r) and interpret the answer.

QUESTION SIX.

[3 + 5 + 2 + 10 marks]

6.1 The following are the numbers of telephone calls received by an automobile rental company in 100 business days.

Number of Telephone Calls	Number of days .
30, but under 35	3
35, but under 40	20
40, but under 45	24
45, but under 50	25
50, but under 55	14
55, but under 60	9
60, but under 65	5

- Compute the mean.
 - Compute the standard deviation.
 - Compute the coefficient of variation and explain what it means.
- 6.2 A motor car manufacturer purchases gear assemblies from a sub-contractor who undertakes to ensure that not more than 5% of his supplies will be defective. In order to provide a check on the quality of incoming supplies a random sample of 200 assemblies is selected of which 17 are found to be defective. Does the sample evidence indicate that the sub-contractor is not maintaining the quality of his supplies at the agreed level? Use $\alpha = 0.01$.

QUESTION SEVEN.

[10 + 10 marks]

In order to determine whether there is a difference in the performance of 2 training methods, samples of individuals from each of the methods were checked. For the 60 individuals from Method A, the mean efficiency score was 35, with a standard deviation of 12. For the 80 individuals from Method B, the mean efficiency score was 27, with a standard deviation of 14.

- Construct a 95% confidence interval of the difference of efficiency scores of these two methods.
- Does the sample evidence indicate any significant difference in efficiency scores of these two methods? Use $\alpha = 0.05$.

Table 1. Binomial Probabilities

Tabulated values are $P(Y \leq a) = \sum_{y=0}^a p(Y)$. (Computations are rounded at third decimal place.)

(a) $n = 5$

a	0.01	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95	0.99
0	.951	.774	.590	.328	.168	.078	.031	.010	.002	.000	.000	.000	.000
1	.999	.977	.919	.737	.528	.337	.188	.087	.031	.007	.000	.000	.000
2	1.000	.999	.991	.942	.837	.683	.500	.317	.163	.038	.009	.001	.000
3	1.000	1.000	1.000	.993	.969	.913	.812	.663	.472	.263	.081	.023	.001
4	1.000	1.000	1.000	1.000	.998	.990	.969	.922	.832	.672	.410	.226	.049

(b) $n = 10$

a	0.01	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95	0.99
0	.904	.599	.349	.107	.028	.006	.001	.000	.000	.000	.000	.000	.000
1	.996	.914	.736	.376	.149	.046	.011	.002	.000	.000	.000	.000	.000
2	1.000	.988	.930	.678	.383	.167	.055	.012	.002	.000	.000	.000	.000
3	1.000	.999	.987	.879	.650	.382	.172	.055	.011	.001	.000	.000	.000
4	1.000	1.000	.998	.967	.850	.633	.377	.166	.047	.006	.000	.000	.000
5	1.000	1.000	1.000	.994	.953	.834	.623	.367	.150	.033	.002	.000	.000
6	1.000	1.000	1.000	.999	.989	.945	.828	.618	.350	.121	.013	.001	.000
7	1.000	1.000	1.000	1.000	.998	.988	.945	.833	.617	.322	.070	.012	.000
8	1.000	1.000	1.000	1.000	1.000	.998	.989	.954	.851	.624	.264	.086	.004
9	1.000	1.000	1.000	1.000	1.000	1.000	.999	.994	.972	.893	.651	.401	.096

(c) $n = 15$

a	0.01	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95	0.99
0	.860	.463	.206	.035	.005	.000	.000	.000	.000	.000	.000	.000	.000
1	.990	.829	.549	.167	.035	.005	.000	.000	.000	.000	.000	.000	.000
2	1.000	.964	.816	.398	.127	.027	.004	.000	.000	.000	.000	.000	.000
3	1.000	.995	.944	.648	.297	.091	.018	.002	.000	.000	.000	.000	.000
4	1.000	.999	.987	.836	.515	.217	.059	.009	.001	.000	.000	.000	.000
5	1.000	1.000	.998	.939	.722	.403	.151	.034	.004	.000	.000	.000	.000
6	1.000	1.000	1.000	.982	.869	.610	.304	.095	.015	.001	.000	.000	.000
7	1.000	1.000	1.000	.996	.950	.787	.500	.213	.050	.004	.000	.000	.000
8	1.000	1.000	1.000	.999	.985	.905	.696	.390	.131	.018	.000	.000	.000
9	1.000	1.000	1.000	1.000	.999	.996	.966	.849	.597	.278	.061	.002	.000
10	1.000	1.000	1.000	1.000	1.000	.999	.991	.941	.783	.485	.164	.013	.001
11	1.000	1.000	1.000	1.000	1.000	1.000	.998	.982	.909	.703	.352	.056	.005
12	1.000	1.000	1.000	1.000	1.000	1.000	.996	.973	.873	.602	.184	.036	.000
13	1.000	1.000	1.000	1.000	1.000	1.000	.995	.995	.965	.833	.451	.171	.010
14	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.995	.995	.965	.794	.537	.140

(d) $n = 20$

a	0.01	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95	0.99
0	.818	.358	.122	.012	.001	.000	.000	.000	.000	.000	.000	.000	.000
1	.983	.736	.392	.069	.008	.001	.000	.000	.000	.000	.000	.000	.000
2	.999	.975	.677	.206	.035	.004	.000	.000	.000	.000	.000	.000	.000
3	1.000	.984	.867	.411	.107	.016	.001	.000	.000	.000	.000	.000	.000
4	1.000	.997	.957	.630	.238	.051	.006	.000	.000	.000	.000	.000	.000
5	1.000	1.000	.989	.804	.416	.126	.021	.002	.000	.000	.000	.000	.000
6	1.000	1.000	.998	.913	.608	.250	.058	.006	.000	.000	.000	.000	.000
7	1.000	1.000	1.000	.968	.772	.416	.132	.021	.001	.000	.000	.000	.000
8	1.000	1.000	1.000	.990	.909	.687	.352	.057	.001	.000	.000	.000	.000
9	1.000	1.000	1.000	.997	.952	.875	.612	.325	.017	.001	.000	.000	.000
10	1.000	1.000	1.000	1.000	.999	.983	.872	.688	.445	.048	.003	.000	.000
11	1.000	1.000	1.000	1.000	1.000	.995	.943	.748	.404	.113	.010	.000	.000
12	1.000	1.000	1.000	1.000	1.000	.999	.979	.868	.584	.228	.032	.000	.000
13	1.000	1.000	1.000	1.000	1.000	1.000	.994	.942	.750	.392	.087	.002	.000
14	1.000	1.000	1.000	1.000	1.000	1.000	.998	.979	.874	.584	.196	.011	.000
15	1.000	1.000	1.000	1.000	1.000	1.000	.994	.949	.762	.370	.043	.003	.000
16	1.000	1.000	1.000	1.000	1.000	1.000	.999	.984	.893	.589	.133	.016	.000
17	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.996	.965	.794	.323	.075	.001
18	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.999	.992	.931	.608	.264	.017
19	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.999	.988	.878	.642	.182	.019

(e) $n = 25$

a	0.01	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95	0.99
0	.778	.277	.072	.004	.000	.000	.000	.000	.000	.000	.000	.000	.000
1	.974	.642	.271	.027	.002	.000	.000	.000	.000	.000	.000	.000	.000
2	.998	.873	.537	.098	.009	.000	.000	.000	.000	.000	.000	.000	.000
3	1.000	.966	.764	.234	.033	.002	.000	.000	.000	.000	.000	.000	.000
4	1.000	.993	.902	.421	.090	.009	.000	.000	.000	.000	.000	.000	.000
5	1.000	.999	.967	.617	.193	.029	.002	.000	.000	.000	.000	.000	.000
6	1.000	1.000	.991	.780	.341	.074	.007	.000	.000	.000	.000	.000	.000
7	1.000	1.000	.998	.891	.512	.154	.022	.001	.000	.000	.000	.000	.000
8	1.000	1.000	1.000	.953	.677	.274	.054	.004	.000	.000	.000	.000	.000
9	1.000	1.000	1.000	.983	.811	.425	.115	.013	.000	.000	.000	.000	.000
10	1.000	1.000	1.000	.994	.902	.586	.212	.034	.002	.000	.000	.000	.000
11	1.000	1.000	1.000	.998	.956	.732	.345	.078	.006	.000	.000	.000	.000
12	1.000	1.000	1.000	1.000	.983	.846	.500	.154	.017	.000	.000	.000	.000
13	1.000	1.000	1.000	1.000	.994	.922	.655	.268	.044	.002	.000	.000	.000
14	1.000	1.000	1.000	1.000	.998	.966	.788	.414	.098	.006	.000	.000	.000
15	1.000	1.000	1.000	1.000	1.000	.987	.885	.575	.189	.017	.000	.000	.000
16	1.000	1.000	1.000	1.000	1.000	.996	.946	.726	.323	.047	.000	.000	.000
17	1.000	1.000	1.000	1.000	1.000	.999	.978	.846	.488	.109	.002	.000	.000
18	1.000	1.000	1.000	1.000	1.000	1.000	.993	.926	.659	.220	.009	.000	.000
19	1.000	1.000	1.000	1.000	1.000	1.000	.998	.971	.807	.383	.033	.001	.000
20	1.000	1.000	1.000	1.000	1.000	1.000	.991	.910	.579	.236	.034	.000	.000
21	1.000	1.000	1.000	1.000	1.000	1.000	.998	.967	.766	.463	.127	.002	.000
22	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.991	.902	.463	.127	.002	.000
23	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.998	.973	.729	.358	.026	.000
24	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.996	.928	.723	.222	.022	.000

Table 2. Table of e^{-x}

x	e^{-x}	x	e^{-x}	x	e^{-x}	x	e^{-x}
0.00	1.000000	2.60	.074274	5.10	.006097	7.60	.000501
0.10	.904837	2.70	.067206	5.20	.005517	7.70	.000453
0.20	.818731	2.80	.060810	5.30	.004992	7.80	.000410
0.30	.740818	2.90	.055023	5.40	.004517	7.90	.000371
0.40	.670320	3.00	.049787	5.50	.004087	8.00	.000336
0.50	.606531	3.10	.045049	5.60	.003698	8.10	.000304
0.60	.548812	3.20	.040762	5.70	.003346	8.20	.000275
0.70	.496585	3.30	.036883	5.80	.003028	8.30	.000249
0.80	.449329	3.40	.033373	5.90	.002739	8.40	.000225
0.90	.406570	3.50	.030197	6.00	.002479	8.50	.000204
1.00	.367879	3.60	.027324	6.10	.002243	8.60	.000184
1.10	.332871	3.70	.024724	6.20	.002029	8.70	.000167
1.20	.301194	3.80	.022371	6.30	.001836	8.80	.000151
1.30	.272532	3.90	.020242	6.40	.001661	8.90	.000136
1.40	.246597	4.00	.018316	6.50	.001503	9.00	.000123
1.50	.223130	4.10	.016573	6.60	.001360	9.10	.000112
1.60	.201897	4.20	.014996	6.70	.001231	9.20	.000101
1.70	.182684	4.30	.013569	6.80	.001114	9.30	.000091
1.80	.165299	4.40	.012277	6.90	.001008	9.40	.000083
1.90	.149569	4.50	.011109	7.00	.000912	9.50	.000075
2.00	.135335	4.60	.010052	7.10	.000825	9.60	.000068
2.10	.122456	4.70	.009095	7.20	.000747	9.70	.000061
2.20	.110803	4.80	.008230	7.30	.000676	9.80	.000056
2.30	.100259	4.90	.007447	7.40	.000611	9.90	.000050
2.40	.090718	5.00	.006738	7.50	.000553	10.00	.000045
2.50	.082085						

Table 3. Poisson Probabilities

$$P(Y \leq d) = \sum_{y=0}^d \frac{e^{-\lambda} \lambda^y}{y!}$$

λ	0	1	2	3	4	5	6	7	8	9
0.02	0.980	1.000	1.000							
0.04	0.961	0.999	1.000							
0.06	0.942	0.998	1.000							
0.08	0.923	0.997	1.000							
0.10	0.905	0.995	1.000							
0.15	0.861	0.990	0.999	1.000						
0.20	0.819	0.982	0.999	1.000						
0.25	0.779	0.974	0.998	1.000						
0.30	0.741	0.963	0.996	1.000						
0.35	0.705	0.951	0.994	1.000						
0.40	0.670	0.938	0.992	0.999	1.000					
0.45	0.638	0.925	0.989	0.999	1.000					
0.50	0.607	0.910	0.986	0.998	1.000					
0.55	0.577	0.894	0.982	0.988	1.000					
0.60	0.549	0.878	0.977	0.997	1.000					
0.65	0.522	0.861	0.972	0.996	0.999	1.000				
0.70	0.497	0.844	0.966	0.994	0.999	1.000				
0.75	0.472	0.827	0.959	0.993	0.999	1.000				
0.80	0.449	0.809	0.953	0.991	0.999	1.000				
0.85	0.427	0.791	0.945	0.989	0.998	1.000				
0.90	0.407	0.772	0.937	0.987	0.998	1.000				
0.95	0.387	0.754	0.929	0.981	0.997	1.000				
1.00	0.368	0.736	0.920	0.981	0.996	0.999	1.000			
1.1	0.333	0.699	0.900	0.974	0.995	0.999	1.000			
1.2	0.301	0.663	0.879	0.966	0.992	0.998	1.000			
1.3	0.273	0.627	0.857	0.957	0.989	0.998	1.000			
1.4	0.247	0.592	0.833	0.946	0.986	0.997	0.999	1.000		
1.5	0.223	0.558	0.809	0.934	0.981	0.996	0.999	1.000		
1.6	0.202	0.525	0.783	0.921	0.976	0.994	0.999	1.000		
1.7	0.183	0.493	0.757	0.907	0.970	0.992	0.998	1.000		
1.8	0.165	0.463	0.731	0.891	0.964	0.990	0.997	0.999	1.000	
1.9	0.150	0.434	0.704	0.875	0.956	0.987	0.997	0.999	1.000	
2.0	0.135	0.406	0.677	0.857	0.947	0.983	0.995	0.999	1.000	


Table 3. (Continued)

λ	0	1	2	3	4	5	6	7	8	9
10.5	0.000	0.000	0.002	0.007	0.021	0.050	0.102	0.179	0.279	0.397
11.0	0.000	0.000	0.001	0.005	0.015	0.038	0.079	0.143	0.232	0.341
11.5	0.000	0.000	0.001	0.003	0.011	0.028	0.060	0.114	0.191	0.289
12.0	0.000	0.000	0.001	0.002	0.008	0.020	0.046	0.090	0.155	0.242
12.5	0.000	0.000	0.000	0.002	0.005	0.015	0.035	0.070	0.125	0.201
13.0	0.000	0.000	0.000	0.001	0.004	0.011	0.026	0.054	0.100	0.166
13.5	0.000	0.000	0.000	0.001	0.003	0.008	0.019	0.041	0.079	0.135
14.0	0.000	0.000	0.000	0.000	0.002	0.006	0.014	0.032	0.062	0.109
14.5	0.000	0.000	0.000	0.000	0.001	0.004	0.010	0.024	0.048	0.088
15.0	0.000	0.000	0.000	0.000	0.001	0.003	0.008	0.018	0.037	0.070
10	10	11	12	13	14	15	16	17	18	19
10.5	0.521	0.639	0.742	0.825	0.888	0.932	0.960	0.978	0.988	0.994
11.0	0.460	0.579	0.689	0.781	0.854	0.907	0.944	0.968	0.982	0.991
11.5	0.402	0.520	0.633	0.733	0.815	0.878	0.924	0.954	0.974	0.986
12.0	0.347	0.462	0.576	0.682	0.772	0.844	0.899	0.937	0.963	0.979
12.5	0.297	0.406	0.519	0.628	0.725	0.806	0.869	0.916	0.948	0.969
13.0	0.252	0.353	0.463	0.573	0.675	0.764	0.835	0.890	0.930	0.957
13.5	0.211	0.304	0.409	0.518	0.623	0.718	0.798	0.861	0.908	0.942
14.0	0.176	0.260	0.358	0.464	0.570	0.669	0.756	0.827	0.883	0.923
14.5	0.145	0.220	0.311	0.413	0.518	0.619	0.711	0.790	0.853	0.901
15.0	0.118	0.185	0.268	0.363	0.466	0.568	0.664	0.749	0.819	0.875
20	20	21	22	23	24	25	26	27	28	29
10.5	0.997	0.999	0.999	1.000						
11.0	0.995	0.998	0.999	1.000						
11.5	0.992	0.996	0.998	0.999	1.000					
12.0	0.988	0.994	0.997	0.999	1.000					
12.5	0.983	0.991	0.995	0.998	0.999	1.000				
13.0	0.975	0.986	0.992	0.996	0.998	0.999	1.000			
13.5	0.965	0.980	0.989	0.994	0.997	0.998	0.999	1.000		
14.0	0.952	0.971	0.983	0.991	0.995	0.997	0.999	0.999	1.000	
14.5	0.936	0.960	0.976	0.986	0.992	0.996	0.998	0.999	1.000	
15.0	0.917	0.947	0.967	0.981	0.989	0.994	0.997	0.998	0.999	1.000

Table 3. (Continued)

λ	4	5	6	7	8	9	10	11	12	13
16	0.000	0.001	0.004	0.010	0.022	0.043	0.077	0.127	0.193	0.275
17	0.000	0.001	0.002	0.005	0.013	0.026	0.049	0.085	0.135	0.201
18	0.000	0.000	0.001	0.003	0.007	0.015	0.030	0.055	0.092	0.143
19	0.000	0.000	0.000	0.001	0.002	0.004	0.009	0.018	0.035	0.061
20	0.000	0.000	0.000	0.000	0.001	0.002	0.005	0.011	0.021	0.039
21	0.000	0.000	0.000	0.000	0.001	0.001	0.003	0.006	0.013	0.025
22	0.000	0.000	0.000	0.000	0.000	0.001	0.002	0.004	0.008	0.015
23	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.002	0.004	0.009
24	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.003	0.005
25	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.001	0.003
14	14	15	16	17	18	19	20	21	22	23
16	0.368	0.467	0.566	0.659	0.742	0.812	0.868	0.911	0.942	0.963
17	0.281	0.371	0.468	0.564	0.655	0.736	0.805	0.861	0.905	0.937
18	0.208	0.287	0.375	0.469	0.562	0.651	0.731	0.799	0.855	0.899
19	0.150	0.215	0.292	0.378	0.469	0.561	0.647	0.725	0.793	0.849
20	0.105	0.157	0.221	0.297	0.381	0.470	0.559	0.644	0.721	0.787
21	0.072	0.111	0.163	0.227	0.302	0.384	0.471	0.558	0.640	0.716
22	0.048	0.077	0.117	0.169	0.232	0.306	0.387	0.472	0.556	0.637
23	0.031	0.052	0.082	0.123	0.175	0.238	0.310	0.389	0.472	0.555
24	0.020	0.034	0.056	0.086	0.128	0.180	0.243	0.314	0.392	0.473
25	0.012	0.022	0.038	0.060	0.092	0.134	0.185	0.247	0.318	0.394
24	24	25	26	27	28	29	30	31	32	33
16	0.978	0.987	0.993	0.996	0.998	0.999	0.999	1.000		
17	0.959	0.975	0.985	0.991	0.995	0.997	0.999	1.000		
18	0.932	0.955	0.972	0.983	0.990	0.994	0.997	0.998	1.000	
19	0.893	0.927	0.951	0.969	0.980	0.988	0.993	0.996	0.998	1.000
20	0.843	0.888	0.922	0.948	0.966	0.978	0.987	0.992	0.995	0.997
21	0.782	0.838	0.883	0.917	0.944	0.963	0.976	0.985	0.991	0.994
22	0.712	0.777	0.832	0.877	0.913	0.940	0.959	0.973	0.983	0.989
23	0.635	0.708	0.772	0.827	0.873	0.908	0.936	0.956	0.971	0.981
24	0.554	0.632	0.704	0.768	0.823	0.868	0.904	0.932	0.953	0.969
25	0.473	0.553	0.629	0.700	0.763	0.818	0.863	0.900	0.929	0.950
34	34	35	36	37	38	39	40	41	42	43
19	0.999	1.000								
20	0.999	0.999	1.000							
21	0.997	0.998	0.999	0.999	1.000					
22	0.994	0.996	0.998	0.999	0.999	1.000				
23	0.988	0.993	0.996	0.997	0.999	0.999	1.000			
24	0.979	0.987	0.992	0.995	0.997	0.998	0.999	1.000		
25	0.966	0.978	0.985	0.991	0.991	0.997	0.998	0.999	1.000	

Table 4. Normal curve areas
Standard normal probability in right-hand tail
(for negative values of z areas are found by symmetry)



z	Second decimal place of z									
	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
2.8	.0026	.0025	.0024	.0023	.0022	.0021	.0021	.0020	.0019	.0019
2.9	.0019	.0018	.0017	.0017	.0016	.0016	.0015	.0015	.0014	.0014
3.0	.00135									
3.5	.000233									
4.0	.0000317									
4.5	.00000340									
5.0	.000000287									

From R. B. Walpole, *Introduction to Statistics* (New York: Macmillan, 1968).

Table 6. Percentage points of the χ^2 -distribution

$\chi^2_{0.100}$	$\chi^2_{0.050}$	$\chi^2_{0.025}$	$\chi^2_{0.010}$	$\chi^2_{0.005}$	d.f.
2.70554	3.84146	5.02389	6.63490	7.87944	1
4.60517	5.99147	7.37776	9.21034	10.59666	2
6.25139	7.81473	9.34840	11.3449	12.8381	3
7.77944	9.48773	11.1433	13.2767	14.8602	4
9.23635	11.0705	12.8325	15.0863	16.7496	5
10.6446	12.5916	14.4494	16.8119	18.5476	6
12.0170	14.0671	16.0128	18.4753	20.2777	7
13.3616	15.5073	17.5346	20.0902	21.9550	8
14.6837	16.9190	19.0228	21.6660	23.5893	9
15.9871	18.3070	20.4831	23.2093	25.1882	10
17.2750	19.6751	21.9200	24.7250	26.7569	11
18.5494	21.0261	23.3367	26.2170	28.2995	12
19.8119	22.3621	24.7356	27.6883	29.8194	13
21.0642	23.6848	26.1190	29.1413	31.3193	14
22.3072	24.9958	27.4884	30.5779	32.8013	15
23.5418	26.2962	28.8454	31.9999	34.2672	16
24.7690	27.5871	30.1910	33.4087	35.7185	17
25.9894	28.8693	31.5264	34.8053	37.1564	18
27.2036	30.1435	32.8523	36.1908	38.5822	19
28.4120	31.4104	34.1696	37.5662	39.9968	20
29.6151	32.6705	35.4789	38.9321	41.4010	21
30.8133	33.9244	36.7807	40.2894	42.7956	22
32.0069	35.1725	38.0757	41.6384	44.1813	23
33.1963	36.4151	39.3641	42.9798	45.5585	24
34.3816	37.6525	40.6465	44.3141	46.9278	25
35.5631	38.8852	41.9232	45.6417	48.2899	26
36.7412	40.1133	43.1944	46.9630	49.6449	27
37.9159	41.3372	44.4607	48.2782	50.9933	28
39.0875	42.5569	45.7222	49.5879	52.3356	29
40.2560	43.7729	46.9792	50.8922	53.6720	30
51.8050	55.7585	59.3417	63.6907	66.7659	40
63.1671	67.5048	71.4202	76.1539	79.4900	50
74.3970	79.0819	83.2976	88.3794	91.9517	60
85.5271	90.5312	95.0231	100.425	104.215	70
96.5782	101.879	106.629	112.329	116.321	80
107.565	113.145	118.136	124.116	128.299	90
118.498	124.342	129.561	135.807	140.169	100

From "Tables of the Percentage Points of the χ^2 -Distribution," *Biometrika*, Vol. 32 (1941), pp. 188-189, by Catherine M. Thompson. Reproduced by permission of Professor H. S. Pearson.