

UNIVERSITY OF ESWATINI
DEPARTMENT OF STATISTICS AND
DEMOGRAPHY
FINAL EXAMINATION PAPER 2019

TITLE OF PAPER : MULTIVARIATE STATISTICS

COURSE CODE : STA 410/ST 410

TIME ALLOWED : 2 HOURS

REQUIREMENTS : STATISTICAL TABLES AND CALCULATOR

INSTRUCTIONS

- 1) Answer any three (3) questions
- 2) Show clearly all your working

THIS PAPER IS NOT TO BE OPENED UNTIL PERMISSION HAS BEEN GRANTED BY THE INVIGILATOR

Question 1

(a) Explain the main purpose of principal component analysis. Briefly discuss the decisions that need to be made when carrying out a principal component analysis. (5)

(b) Consider the following data;

| X_1 | X_2 |
|-------|-------|
| 1.00 | 3.00 |
| 1.00 | 2.00 |
| 2.00 | 3.00 |
| 0.00 | 3.00 |
| 5.00 | 4.00 |
| 4.00 | 5.00 |
| 5.00 | 5.00 |
| 3.00 | 4.00 |

- i. Calculate the covariance matrix for this data. (3)
- ii. Determine the principal components and find the proportion of the total variance explained by the first component. (12)

Question 2

A certain type of tropical disease is characterized by fever, low blood pressure and body aches. A pharmaceutical company is working on a new drug to treat this type of disease and wanted to determine whether the drug is effective. They took a random sample of 20 people with this type of disease and 18 with a placebo. Based on the data below determine whether the drug is effective at reducing these two symptoms. State the assumptions for your test. (20)

| Drug | | Placebo | |
|-------|----------|---------|----------|
| Fever | Pressure | Fever | Pressure |
| 38.4 | 73 | 40.9 | 54 |
| 36.8 | 85 | 39.5 | 75 |
| 40.0 | 58 | 39.4 | 57 |
| 39.8 | 80 | 38.2 | 71 |
| 38.6 | 68 | 39.7 | 65 |
| 39.1 | 52 | 38.9 | 49 |
| 38.9 | 79 | 38.6 | 58 |
| 36.8 | 100 | 39.9 | 52 |
| 40.4 | 64 | 41.3 | 62 |
| 39.4 | 53 | 38.1 | 57 |
| 38.0 | 70 | 39.6 | 78 |
| 38.6 | 75 | 37.1 | 92 |
| 40.1 | 48 | 39.5 | 63 |
| 38.1 | 57 | 40.3 | 52 |
| 37.2 | 78 | 41.5 | 46 |
| 39.5 | 65 | 39.3 | 56 |
| 37.3 | 77 | 37.6 | 86 |
| 39.1 | 67 | 40.6 | 48 |
| 39.9 | 52 | | |
| 37.8 | 68 | | |

Question 3

- (a) Briefly describe the purposes of cluster analysis. (3)
- (b) Suppose that objects are to be clustered on the basis of an observation vector that consists of p continuous measurements. Define the Euclidean distance between objects with observation vectors

$$\mathbf{x} = [x_1 \ x_2 \ \dots \ x_p]^T, \text{ and } \mathbf{y} = [y_1 \ y_2 \ \dots \ y_p]^T.$$

- Also define any two of the following alternative similarity measures: squared Euclidean distance; Manhattan (or city-block) distance; Chebyshev distance; Mahalanobis distance. For each of your chosen cases, describe how the overall features of the complete set of distances obtained using that measure would differ from the corresponding set of Euclidean distances. (5)
- (c) Describe the single-linkage and complete-linkage approaches to hierarchical clustering. How might the outcome of a cluster analysis differ depending on which of these approaches was used? (6)
- (d) Describe k -means clustering. Mention one advantage and one disadvantage of this method compared with hierarchical procedures. (6)

Question 4

Conn's Syndrome is a form of hypertension that has two possible causes: an adenoma (Type A patient), which has to be removed by surgery, and bilateral hyperplasia (Type B patient), which is a more diffuse condition and is treated with drugs. It can be hard to tell whether a patient is Type A or Type B. Researchers investigated a group of 31 sufferers of Conn's Syndrome, recording their age (in years) and the concentrations of the following three chemicals in blood plasma (in meq/l): sodium, potassium, carbon dioxide. All these patients then underwent surgery, which revealed that 20 of them were Type A and the other 11 Type B. The computer output displayed on the next page is from an analysis of the data for all 31 patients in the study.

- i. What does the matrix plot suggest about the potential usefulness of each of these variables for classifying patients as Type A or Type B? (2)
- ii. A linear discriminant analysis of the data was carried out, with the results shown in Table 1. Explain what is meant by leaving-one-out cross-validation. Give one reason for carrying out this procedure as part of discriminant analysis. What do the results in Table 1 suggest about the usefulness of the linear discriminant?
- iii. The linear discriminant function is

$$y = -20.0 - (0.1 \times \text{age}) + (0.2 \times \text{sodium}) - (3.0 \times \text{potassium}) + (0.6 \times \text{carbon dioxide})$$

, where positive values of y indicate Type A patients. Use it to classify another Conn's Syndrome sufferer, who is 40 years old and has the following test results: sodium 144.1, potassium 3.4, carbon dioxide 25.2 meq/l. (2)

- iv. List the main statistical assumptions required to justify the use of linear discrimination. In what way may these assumptions be relaxed if quadratic discrimination is used instead of linear discrimination? (4)
- v. Table 2 gives the results of quadratic discrimination, in a form comparable to Table 1. Compare the results from the two discriminants. Which would you recommend using in practice, and why? (4)

Matrix Plot




Table 1: Results of Linear Discriminant Analysis

(a) Without Cross-Validation

| Put into Group: | True Group | |
|-----------------|------------|-------|
| | A | B |
| A | 17 | 0 |
| B | 2 | 11 |
| Total | 20 | 11 |
| No. correct | 17 | 11 |
| Proportion | 0.850 | 1.000 |

Overall Proportion Correct = 0.903

(b) With Cross-Validation

| Put into Group: | True Group | |
|-----------------|------------|-------|
| | A | B |
| A | 16 | 1 |
| B | 4 | 10 |
| Total | 20 | 11 |
| No. correct | 16 | 10 |
| Proportion | 0.800 | 0.909 |

Overall Proportion Correct = 0.839

Table 2: Results of Quadratic Discriminant Analysis

(a) Without Cross-Validation

| Put into Group: | True Group | |
|-----------------|------------|-------|
| | A | B |
| A | 19 | 0 |
| B | 1 | 11 |
| Total | 20 | 11 |
| No. correct | 19 | 11 |
| Proportion | 0.950 | 1.000 |


Overall Proportion Correct = 0.968

(b) With Cross-Validation

| Put into Group: | True Group | |
|-----------------|------------|-------|
| | A | B |
| A | 17 | 3 |
| B | 3 | 8 |
| Total | 20 | 11 |
| No. correct | 17 | 8 |
| Proportion | 0.850 | 0.727 |


Overall Proportion Correct = 0.806

Table 4 Normal Curve Areas
 Standard normal probability in right-hand tail
 (for negative values of z , areas are found by symmetry)



| z | Second decimal place of z | | | | | | | | | |
|-----|-----------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | .00 | .01 | .02 | .03 | .04 | .05 | .06 | .07 | .08 | .09 |
| 0.0 | .5000 | .4960 | .4920 | .4880 | .4840 | .4801 | .4761 | .4721 | .4681 | .4641 |
| 0.1 | .4602 | .4562 | .4522 | .4483 | .4443 | .4404 | .4364 | .4325 | .4286 | .4247 |
| 0.2 | .4207 | .4168 | .4129 | .4090 | .4052 | .4013 | .3974 | .3936 | .3897 | .3859 |
| 0.3 | .3821 | .3783 | .3745 | .3707 | .3669 | .3632 | .3594 | .3557 | .3520 | .3483 |
| 0.4 | .3446 | .3409 | .3372 | .3336 | .3300 | .3264 | .3228 | .3192 | .3156 | .3121 |
| 0.5 | .3085 | .3050 | .3015 | .2981 | .2946 | .2912 | .2877 | .2843 | .2810 | .2776 |
| 0.6 | .2743 | .2709 | .2676 | .2643 | .2611 | .2578 | .2546 | .2514 | .2483 | .2451 |
| 0.7 | .2420 | .2389 | .2358 | .2327 | .2296 | .2266 | .2236 | .2206 | .2177 | .2148 |
| 0.8 | .2119 | .2090 | .2061 | .2033 | .2005 | .1977 | .1949 | .1922 | .1894 | .1867 |
| 0.9 | .1841 | .1814 | .1788 | .1762 | .1736 | .1711 | .1685 | .1660 | .1635 | .1611 |
| 1.0 | .1587 | .1562 | .1539 | .1515 | .1492 | .1469 | .1446 | .1423 | .1401 | .1379 |
| 1.1 | .1357 | .1335 | .1314 | .1292 | .1271 | .1251 | .1230 | .1210 | .1190 | .1170 |
| 1.2 | .1151 | .1131 | .1112 | .1093 | .1075 | .1056 | .1038 | .1020 | .1003 | .0985 |
| 1.3 | .0968 | .0951 | .0934 | .0918 | .0901 | .0885 | .0869 | .0853 | .0838 | .0823 |
| 1.4 | .0808 | .0793 | .0778 | .0764 | .0749 | .0735 | .0722 | .0708 | .0694 | .0681 |
| 1.5 | .0668 | .0655 | .0643 | .0630 | .0618 | .0606 | .0594 | .0582 | .0571 | .0559 |
| 1.6 | .0548 | .0537 | .0526 | .0516 | .0505 | .0495 | .0485 | .0475 | .0465 | .0455 |
| 1.7 | .0446 | .0436 | .0427 | .0418 | .0409 | .0401 | .0392 | .0384 | .0375 | .0367 |
| 1.8 | .0359 | .0352 | .0344 | .0336 | .0329 | .0322 | .0314 | .0307 | .0301 | .0294 |
| 1.9 | .0287 | .0281 | .0274 | .0268 | .0262 | .0256 | .0250 | .0244 | .0239 | .0233 |
| 2.0 | .0228 | .0222 | .0217 | .0212 | .0207 | .0202 | .0197 | .0192 | .0188 | .0183 |
| 2.1 | .0179 | .0174 | .0170 | .0166 | .0162 | .0158 | .0154 | .0150 | .0146 | .0143 |
| 2.2 | .0139 | .0136 | .0132 | .0129 | .0125 | .0122 | .0119 | .0116 | .0113 | .0110 |
| 2.3 | .0107 | .0104 | .0102 | .0099 | .0096 | .0094 | .0091 | .0089 | .0087 | .0084 |
| 2.4 | .0082 | .0080 | .0078 | .0075 | .0073 | .0071 | .0069 | .0068 | .0066 | .0064 |
| 2.5 | .0062 | .0060 | .0059 | .0057 | .0055 | .0054 | .0052 | .0051 | .0049 | .0048 |
| 2.6 | .0047 | .0045 | .0044 | .0043 | .0041 | .0040 | .0039 | .0038 | .0037 | .0036 |
| 2.7 | .0035 | .0034 | .0033 | .0032 | .0031 | .0030 | .0029 | .0028 | .0027 | .0026 |
| 2.8 | .0026 | .0025 | .0024 | .0023 | .0023 | .0022 | .0021 | .0021 | .0020 | .0019 |
| 2.9 | .0019 | .0018 | .0017 | .0017 | .0016 | .0016 | .0015 | .0015 | .0014 | .0014 |
| 3.0 | .00135 | | | | | | | | | |
| 3.5 | .000 233 | | | | | | | | | |
| 4.0 | .000 031 7 | | | | | | | | | |
| 4.5 | .000 003 40 | | | | | | | | | |
| 5.0 | .000 000 287 | | | | | | | | | |

From R. E. Walpole, *Introduction to Statistics* (New York: Macmillan, 1968).

Table 6 Percentage Points of the χ^2 Distributions


| df | $\chi^2_{0.995}$ | $\chi^2_{0.990}$ | $\chi^2_{0.975}$ | $\chi^2_{0.950}$ | $\chi^2_{0.900}$ |
|-----|------------------|------------------|------------------|------------------|------------------|
| 1 | 0.0000393 | 0.0001571 | 0.0009821 | 0.0039321 | 0.0157908 |
| 2 | 0.0100251 | 0.0201007 | 0.0506356 | 0.102587 | 0.210720 |
| 3 | 0.0717212 | 0.114832 | 0.215795 | 0.351846 | 0.584375 |
| 4 | 0.206990 | 0.297110 | 0.484419 | 0.710721 | 1.063623 |
| 5 | 0.411740 | 0.554300 | 0.831211 | 1.145476 | 1.61031 |
| 6 | 0.675727 | 0.872085 | 1.237347 | 1.63539 | 2.20413 |
| 7 | 0.989265 | 1.239043 | 1.68987 | 2.16735 | 2.83311 |
| 8 | 1.344419 | 1.646482 | 2.17973 | 2.73264 | 3.48954 |
| 9 | 1.734926 | 2.087912 | 2.70039 | 3.32511 | 4.16816 |
| 10 | 2.15585 | 2.55821 | 3.24697 | 3.94030 | 4.86518 |
| 11 | 2.60321 | 3.05347 | 3.81575 | 4.57481 | 5.57779 |
| 12 | 3.07382 | 3.57056 | 4.40379 | 5.22603 | 6.30380 |
| 13 | 3.56503 | 4.10691 | 5.00874 | 5.89186 | 7.04150 |
| 14 | 4.07468 | 4.66043 | 5.62872 | 6.57063 | 7.78953 |
| 15 | 4.60094 | 5.22935 | 6.26214 | 7.26094 | 8.54675 |
| 16 | 5.14224 | 5.81221 | 6.90766 | 7.96164 | 9.31223 |
| 17 | 5.69724 | 6.40776 | 7.56418 | 8.67176 | 10.0852 |
| 18 | 6.26481 | 7.01491 | 8.23075 | 9.39046 | 10.8649 |
| 19 | 6.84398 | 7.63273 | 8.90655 | 10.1170 | 11.6509 |
| 20 | 7.43386 | 8.26040 | 9.59083 | 10.8508 | 12.4426 |
| 21 | 8.03366 | 8.89720 | 10.28293 | 11.5913 | 13.2396 |
| 22 | 8.64272 | 9.54249 | 10.9823 | 12.3380 | 14.0415 |
| 23 | 9.26042 | 10.19567 | 11.6885 | 13.0905 | 14.8479 |
| 24 | 9.88623 | 10.8564 | 12.4011 | 13.8484 | 15.6587 |
| 25 | 10.5197 | 11.5240 | 13.1197 | 14.6114 | 16.4734 |
| 26 | 11.1603 | 12.1981 | 13.8439 | 15.3791 | 17.2919 |
| 27 | 11.8076 | 12.8786 | 14.5733 | 16.1513 | 18.1138 |
| 28 | 12.4613 | 13.5648 | 15.3079 | 16.9279 | 18.9392 |
| 29 | 13.1211 | 14.2565 | 16.0471 | 17.7083 | 19.7677 |
| 30 | 13.7867 | 14.9535 | 16.7908 | 18.4926 | 20.5992 |
| 40 | 20.7065 | 22.1643 | 24.4331 | 26.5093 | 29.0505 |
| 50 | 27.9907 | 29.7067 | 32.3574 | 34.7642 | 37.6886 |
| 60 | 35.5346 | 37.4848 | 40.4817 | 43.1879 | 46.4589 |
| 70 | 43.2752 | 45.4418 | 48.7576 | 51.7393 | 55.3290 |
| 80 | 51.1720 | 53.5400 | 57.1532 | 60.3915 | 64.2778 |
| 90 | 59.1963 | 61.7541 | 65.6466 | 69.1260 | 73.2912 |
| 100 | 67.3276 | 70.0648 | 74.2219 | 77.9295 | 82.3581 |

Table 6 (Continued)

| $\chi^2_{0.100}$ | $\chi^2_{0.050}$ | $\chi^2_{0.025}$ | $\chi^2_{0.010}$ | $\chi^2_{0.005}$ | df |
|------------------|------------------|------------------|------------------|------------------|-----|
| 2.70554 | 3.84146 | 5.02389 | 6.63490 | 7.87944 | 1 |
| 4.60517 | 5.99147 | 7.37776 | 9.21034 | 10.5966 | 2 |
| 6.25139 | 7.81473 | 9.34840 | 11.3449 | 12.8381 | 3 |
| 7.77944 | 9.48773 | 11.1433 | 13.2767 | 14.8602 | 4 |
| 9.23635 | 11.0705 | 12.8325 | 15.0863 | 16.7496 | 5 |
| 10.6446 | 12.5916 | 14.4494 | 16.8119 | 18.5476 | 6 |
| 12.0170 | 14.0671 | 16.0128 | 18.4753 | 20.2777 | 7 |
| 13.3616 | 15.5073 | 17.5346 | 20.0902 | 21.9550 | 8 |
| 14.6837 | 16.9190 | 19.0228 | 21.6660 | 23.5893 | 9 |
| 15.9871 | 18.3070 | 20.4831 | 23.2093 | 25.1882 | 10 |
| 17.2750 | 19.6751 | 21.9200 | 24.7250 | 26.7569 | 11 |
| 18.5494 | 21.0261 | 23.3367 | 26.2170 | 28.2995 | 12 |
| 19.8119 | 22.3621 | 24.7356 | 27.6883 | 29.8194 | 13 |
| 21.0642 | 23.6848 | 26.1190 | 29.1413 | 31.3193 | 14 |
| 22.3072 | 24.9958 | 27.4884 | 30.5779 | 32.8013 | 15 |
| 23.5418 | 26.2962 | 28.8454 | 31.9999 | 34.2672 | 16 |
| 24.7690 | 27.5871 | 30.1910 | 33.4087 | 35.7185 | 17 |
| 25.9894 | 28.8693 | 31.5264 | 34.8053 | 37.1564 | 18 |
| 27.2036 | 30.1435 | 32.8523 | 36.1908 | 38.5822 | 19 |
| 28.4120 | 31.4104 | 34.1696 | 37.5662 | 39.9968 | 20 |
| 29.6151 | 32.6705 | 35.4789 | 38.9321 | 41.4010 | 21 |
| 30.8133 | 33.9244 | 36.7807 | 40.2894 | 42.7956 | 22 |
| 32.0069 | 35.1725 | 38.0757 | 41.6384 | 44.1813 | 23 |
| 33.1963 | 36.4151 | 39.3641 | 42.9798 | 45.5585 | 24 |
| 34.3816 | 37.6525 | 40.6465 | 44.3141 | 46.9278 | 25 |
| 35.5631 | 38.8852 | 41.9232 | 45.6417 | 48.2899 | 26 |
| 36.7412 | 40.1133 | 43.1944 | 46.9630 | 49.6449 | 27 |
| 37.9159 | 41.3372 | 44.4607 | 48.2782 | 50.9933 | 28 |
| 39.0875 | 42.5569 | 45.7222 | 49.5879 | 52.3356 | 29 |
| 40.2560 | 43.7729 | 46.9792 | 50.8922 | 53.6720 | 30 |
| 51.8050 | 55.7585 | 59.3417 | 63.6907 | 66.7659 | 40 |
| 63.1671 | 67.5048 | 71.4202 | 76.1539 | 79.4900 | 50 |
| 74.3970 | 79.0819 | 83.2976 | 88.3794 | 91.9517 | 60 |
| 85.5271 | 90.5312 | 95.0231 | 100.425 | 104.215 | 70 |
| 96.5782 | 101.879 | 106.629 | 112.329 | 116.321 | 80 |
| 107.565 | 113.145 | 118.136 | 124.116 | 128.299 | 90 |
| 118.498 | 124.342 | 129.561 | 135.807 | 140.169 | 100 |

From "Tables of the Percentage Points of the χ^2 -Distribution." *Biometrika*, Vol. 32 (1941), pp. 188-189, by Catherine M. Thompson.

Table 7 Percentage Points of the F Distributions



| Denominator df | Numerator df | | | | | | | | | |
|----------------|--------------|-------|--------|-------|-------|-------|-------|-------|-------|-------|
| | α | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 1 | .100 | 39.86 | 49.50 | 53.59 | 55.83 | 57.24 | 58.20 | 58.91 | 59.44 | 59.86 |
| | .050 | 161.4 | 199.5 | 215.7 | 224.6 | 230.2 | 234.0 | 236.8 | 238.9 | 240.5 |
| | .025 | 647.8 | 799.5 | 864.2 | 899.6 | 921.8 | 937.1 | 948.2 | 956.7 | 963.3 |
| | .010 | 4052 | 4999.5 | 5403 | 5625 | 5764 | 5859 | 5928 | 5982 | 6022 |
| | .005 | 16211 | 20000 | 21615 | 22500 | 23056 | 23437 | 23715 | 23925 | 24091 |
| 2 | .100 | 8.53 | 9.00 | 9.16 | 9.24 | 9.29 | 9.33 | 9.35 | 9.37 | 9.38 |
| | .050 | 18.51 | 19.00 | 19.16 | 19.25 | 19.30 | 19.33 | 19.35 | 19.37 | 19.38 |
| | .025 | 38.51 | 39.00 | 39.17 | 39.25 | 39.30 | 39.33 | 39.36 | 39.37 | 39.39 |
| | .010 | 98.50 | 99.00 | 99.17 | 99.25 | 99.30 | 99.33 | 99.36 | 99.37 | 99.39 |
| | .005 | 198.5 | 199.0 | 199.2 | 199.2 | 199.3 | 199.3 | 199.4 | 199.4 | 199.4 |
| 3 | .100 | 5.54 | 5.46 | 5.39 | 5.34 | 5.31 | 5.28 | 5.27 | 5.25 | 5.24 |
| | .050 | 10.13 | 9.55 | 9.28 | 9.12 | 9.01 | 8.94 | 8.89 | 8.85 | 8.81 |
| | .025 | 17.44 | 16.04 | 15.44 | 15.10 | 14.88 | 14.73 | 14.62 | 14.54 | 14.47 |
| | .010 | 34.12 | 30.82 | 29.46 | 28.71 | 28.24 | 27.91 | 27.67 | 27.49 | 27.35 |
| | .005 | 55.55 | 49.80 | 47.47 | 46.19 | 45.39 | 44.84 | 44.43 | 44.13 | 43.88 |
| 4 | .100 | 4.54 | 4.32 | 4.19 | 4.11 | 4.05 | 4.01 | 3.98 | 3.95 | 3.94 |
| | .050 | 7.71 | 6.94 | 6.59 | 6.39 | 6.26 | 6.16 | 6.09 | 6.04 | 6.00 |
| | .025 | 12.22 | 10.65 | 9.98 | 9.60 | 9.36 | 9.20 | 9.07 | 8.98 | 8.90 |
| | .010 | 21.20 | 18.00 | 16.69 | 15.98 | 15.52 | 15.21 | 14.98 | 14.80 | 14.66 |
| | .005 | 31.33 | 26.28 | 24.26 | 23.15 | 22.46 | 21.97 | 21.62 | 21.35 | 21.14 |
| 5 | .100 | 4.06 | 3.78 | 3.62 | 3.52 | 3.45 | 3.40 | 3.37 | 3.34 | 3.32 |
| | .050 | 6.61 | 5.79 | 5.41 | 5.19 | 5.05 | 4.95 | 4.88 | 4.82 | 4.77 |
| | .025 | 10.01 | 8.43 | 7.76 | 7.39 | 7.15 | 6.98 | 6.85 | 6.76 | 6.68 |
| | .010 | 16.26 | 13.27 | 12.06 | 11.39 | 10.97 | 10.67 | 10.46 | 10.29 | 10.16 |
| | .005 | 22.78 | 18.31 | 16.53 | 15.56 | 14.94 | 14.51 | 14.20 | 13.96 | 13.77 |
| 6 | .100 | 3.78 | 3.46 | 3.29 | 3.18 | 3.11 | 3.05 | 3.01 | 2.98 | 2.96 |
| | .050 | 5.99 | 5.14 | 4.76 | 4.53 | 4.39 | 4.28 | 4.21 | 4.15 | 4.10 |
| | .025 | 8.81 | 7.26 | 6.60 | 6.23 | 5.99 | 5.82 | 5.70 | 5.60 | 5.52 |
| | .010 | 13.75 | 10.92 | 9.78 | 9.15 | 8.75 | 8.47 | 8.26 | 8.10 | 7.98 |
| | .005 | 18.63 | 14.54 | 12.92 | 12.03 | 11.46 | 11.07 | 10.79 | 10.57 | 10.39 |
| 7 | .100 | 3.59 | 3.26 | 3.07 | 2.96 | 2.88 | 2.83 | 2.78 | 2.75 | 2.72 |
| | .050 | 5.59 | 4.74 | 4.35 | 4.12 | 3.97 | 3.87 | 3.79 | 3.73 | 3.68 |
| | .025 | 8.07 | 6.54 | 5.89 | 5.52 | 5.29 | 5.12 | 4.99 | 4.90 | 4.82 |
| | .010 | 12.25 | 9.55 | 8.45 | 7.85 | 7.46 | 7.19 | 6.99 | 6.84 | 6.72 |
| | .005 | 16.24 | 12.40 | 10.88 | 10.05 | 9.52 | 9.16 | 8.89 | 8.68 | 8.51 |

Table 7 (Continued)

| Denominator df | F_{α} | | | | | | | | | |
|-------------------|--------------|------|------|------|------|------|------|------|------|------|
| | α | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 29 | .100 | 2.89 | 2.50 | 2.28 | 2.15 | 2.06 | 1.99 | 1.93 | 1.89 | 1.86 |
| | .050 | 4.18 | 3.33 | 2.93 | 2.70 | 2.55 | 2.43 | 2.35 | 2.28 | 2.22 |
| | .025 | 5.59 | 4.20 | 3.61 | 3.27 | 3.04 | 2.88 | 2.76 | 2.67 | 2.59 |
| | .010 | 7.60 | 5.42 | 4.54 | 4.04 | 3.73 | 3.50 | 3.33 | 3.20 | 3.09 |
| | .005 | 9.23 | 6.40 | 5.28 | 4.66 | 4.26 | 3.98 | 3.77 | 3.61 | 3.48 |
| 30 | .100 | 2.88 | 2.49 | 2.28 | 2.14 | 2.05 | 1.98 | 1.93 | 1.88 | 1.85 |
| | .050 | 4.17 | 3.32 | 2.92 | 2.69 | 2.53 | 2.42 | 2.33 | 2.27 | 2.21 |
| | .025 | 5.57 | 4.18 | 3.59 | 3.25 | 3.03 | 2.87 | 2.75 | 2.65 | 2.57 |
| | .010 | 7.56 | 5.39 | 4.51 | 4.02 | 3.70 | 3.47 | 3.30 | 3.17 | 3.07 |
| | .005 | 9.18 | 6.35 | 5.24 | 4.62 | 4.23 | 3.95 | 3.74 | 3.58 | 3.45 |
| 40 | .100 | 2.84 | 2.44 | 2.23 | 2.09 | 2.00 | 1.93 | 1.87 | 1.83 | 1.79 |
| | .050 | 4.08 | 3.23 | 2.84 | 2.61 | 2.45 | 2.34 | 2.25 | 2.18 | 2.12 |
| | .025 | 5.42 | 4.05 | 3.46 | 3.13 | 2.90 | 2.74 | 2.62 | 2.53 | 2.45 |
| | .010 | 7.31 | 5.18 | 4.31 | 3.83 | 3.51 | 3.29 | 3.12 | 2.99 | 2.89 |
| | .005 | 8.83 | 6.07 | 4.98 | 4.37 | 3.99 | 3.71 | 3.51 | 3.35 | 3.22 |
| 60 | .100 | 2.79 | 2.39 | 2.18 | 2.04 | 1.95 | 1.87 | 1.82 | 1.77 | 1.74 |
| | .050 | 4.00 | 3.15 | 2.76 | 2.53 | 2.37 | 2.25 | 2.17 | 2.10 | 2.04 |
| | .025 | 5.29 | 3.93 | 3.34 | 3.01 | 2.79 | 2.63 | 2.51 | 2.41 | 2.33 |
| | .010 | 7.08 | 4.98 | 4.13 | 3.65 | 3.34 | 3.12 | 2.95 | 2.82 | 2.72 |
| | .005 | 8.49 | 5.79 | 4.73 | 4.14 | 3.76 | 3.49 | 3.29 | 3.13 | 3.01 |
| 120 | .100 | 2.75 | 2.35 | 2.13 | 1.99 | 1.90 | 1.82 | 1.77 | 1.72 | 1.68 |
| | .050 | 3.92 | 3.07 | 2.68 | 2.45 | 2.29 | 2.17 | 2.09 | 2.02 | 1.96 |
| | .025 | 5.15 | 3.80 | 3.23 | 2.89 | 2.67 | 2.52 | 2.39 | 2.30 | 2.22 |
| | .010 | 6.85 | 4.79 | 3.95 | 3.48 | 3.17 | 2.96 | 2.79 | 2.66 | 2.56 |
| | .005 | 8.18 | 5.54 | 4.50 | 3.92 | 3.55 | 3.28 | 3.09 | 2.93 | 2.81 |
| ∞ | .100 | 2.71 | 2.30 | 2.08 | 1.94 | 1.85 | 1.77 | 1.72 | 1.67 | 1.63 |
| | .050 | 3.84 | 3.00 | 2.60 | 2.37 | 2.21 | 2.10 | 2.01 | 1.94 | 1.88 |
| | .025 | 5.02 | 3.69 | 3.12 | 2.79 | 2.57 | 2.41 | 2.29 | 2.19 | 2.11 |
| | .010 | 6.63 | 4.61 | 3.78 | 3.32 | 3.02 | 2.80 | 2.64 | 2.51 | 2.41 |
| | .005 | 7.88 | 5.30 | 4.28 | 3.72 | 3.35 | 3.09 | 2.90 | 2.74 | 2.62 |

From "Tables of percentage points of the inverted beta (F) distribution." *Biometrika*, Vol. 33 (1943) by M. Merrington and C. M. Thompson and from Table 18 of *Biometrika Tables for Statisticians*, Vol. 1, Cambridge University Press, 1954, edited by E. S. Pearson and H. O. Hartley.